arXiv:1507.02312v3 [math.SP] 9 Apr 2017

Extension theory approach in stability of standing waves for NLS equation with point interactions

Jaime Angulo Pava and Nataly Goloshchapova

Abstract

The aim of this work is to demonstrate the effectiveness of the extension theory for the investigation of the stability of standing waves for semilinear Schrödinger equations with δ- and δ'- interaction on the line and on the star graph.

Mathematics Subject Classification (2000). Primary 35Q51, 35J61; Secondary 47E05.
Key words. Nonlinear Schrödinger equation, point interaction, self-adjoint extension, deficiency indices, semi-bounded operator, Sturm-Liouville theory, orbital stability, standing wave, star graph.

1 Introduction

In the last years the study of nonlinear dispersive models with point interactions has attracted a lot of attention of mathematicians and physicists. In particular, such models appear in nonlinear optics, Bose Einstein condensates, and quantum graphs (or networks) (see [13, 17, 18, 31] and reference therein). The prototype equation for description of these models is the nonlinear Schrödinger equation (NLS henceforth)

$$i\partial_t u(t,x) + \partial_x^2 u(t,x) = \mu\,|u(t,x)|^{p-1}\,u(t,x), \quad x \neq 0, \quad (t,x) \in \mathbb{R}\times\mathbb{R}, \quad p > 1, \tag{1.1}$$

with specific boundary conditions at $x = 0$ induced by the certain impurity. Recently, there has been a growing interest in this model from the mathematical side in the attempt to establish the local well posedness of its initial value problem and periodic boundary value problem, the long time behavior of its solutions, blow up and scattering results, the existence of stationary states (see [1, 10–12, 18, 19, 22, 25, 28] and reference therein).

Numerous analytical, numerical and experimental works deal with special solutions of (1.1). In particular, a big part of them consider so-called *standing wave solutions* which preserve the spatial shape and harmonically oscillate in time, namely, solutions of the form

$$u(t,x) = e^{i\omega t}\varphi_\omega(x).$$

Both the structure of the family of standing waves, as well as the problems of the existence and stability/instability, are far richer and more complicated in the case of the NLS equation with point interactions than in the case of the NLS equation without impurities.

In this paper we investigate an orbital stability of the standing waves of the NLS equation with δ- and δ'-interactions on the line and on the star graph. One of the main advantages of using the point interactions is the existence of an explicit expression for the soliton profile φ_ω. This allows one to prove very specific results, the proofs of which are considerably harder in the case of general linear potential.

Our paper consists of two parts. In the first part we propose a new approach to the known stability results for NLS equation with δ- and δ'-interaction on the line. It should be noted that most of previous results are based on either variational methods or the abstract stability theory by Grillakis, Shatah and Strauss [26, 27] which requires spectral analysis of certain self-adjoint Schrödinger operators. In particular, investigation of the spectrum is based on analytic perturbation theory and variational methods.

Our approach relies on the theory of extensions of symmetric operators, Sturm oscillation theorem, and spectral theory of self-adjoint Schrödinger operators. Our method simplifies essentially the stability analysis at least in the case of attractive point interactions.

We also show that stability of standing wave solution of NLS-δ equation with repulsive nonlinearity ($\mu > 0$ in (1.1)) might be easily established, while in [30] the authors used variational approach which requires considerably more extensive proofs. It is worth mentioning that extension theory was applied in [11] for investigation of stability of standing waves with the peak-Gausson profile $\varphi_{\omega,\gamma}(x) = e^{\frac{\omega+1}{2}} e^{-\frac{1}{2}(|x|+\frac{\gamma}{2})^2}$ of the following NLS-log-δ equation

$$i\partial_t u + \partial_x^2 u - \gamma\delta(x)u + u\mathrm{Log}|u|^2 = 0, \quad x \neq 0, \quad \gamma > 0.$$

The second part of the paper is devoted to the study of the orbital stability of the NLS equation with δ- and δ'-interaction on the star graph. To our knowledge such type of models is much more less studied. For instance, the only stability results for NLS-δ equation on the graph were obtained in [1, 2], meanwhile the investigation for NLS-δ' equation on the graph have never been elaborated before. Here we propose a short proof of the orbital stability of a certain family of standing waves for NLS-δ equation on the graph with negative intensity $\alpha < 0$. Note that in [1] it was a big problem to show the stability for any $\alpha < 0$ without restriction $\alpha < \alpha^* < 0$. Our approach allows us to overcame easily this difficulty. Moreover, in the present work we prove the following novel stability theorem for the standing waves of NLS-δ' equation on the graph with specific profile.

Theorem 1.1. *Let* $\lambda < 0$, $1 < p \le 5$, *and* $\omega > \frac{N^2}{\lambda^2}$. *Let also* $\mathbf{U}(t,x) = e^{it\omega}\mathbf{\Phi}_0^\lambda = (e^{it\omega}\varphi_0^\lambda)_{j=1}^N$ *be the standing wave of* (2.11) *defined by* (2.20). *Then the following assertions hold.*

(*i*) *If* $\omega < \frac{N^2}{\lambda^2}\frac{p+1}{p-1}$, *then* $\mathbf{U}(t,x)$ *is orbitally stable in* $H^1(\mathcal{G})$.

(*ii*) *If* $\omega > \frac{N^2}{\lambda^2}\frac{p+1}{p-1}$ *and* N *is even, then* $\mathbf{U}(t,x)$ *is orbitally unstable in* $H^1(\mathcal{G})$.

The above result partially generalizes the results for the NLS-δ' equation on the line.

The paper is organized as follows. In the Preliminaries we give some brief description of the point interactions in the context of the theory of extensions and also review previous results on the orbital stability. In Section 3 we discuss NLS equation with δ- and δ'-interaction on the line, while in Section 4 we provide investigation on the star graph.

Notation. Let $-\infty \le a < b \le \infty$. We denote by $L^2(a,b)$ the Hilbert space equipped with the inner product $(u,v) = \mathrm{Re}\int\limits_a^b u(x)\overline{v(x)}dx$. By $H^1(\mathbb{R})$, $H^2(\mathbb{R}\setminus\{0\}) = H^2(\mathbb{R}_-) \oplus H^2(\mathbb{R}_+)$

we denote the Sobolev spaces. The space of radial symmetric functions in $H^1(\mathbb{R})$ is denoted by $H^1_{\text{rad}}(\mathbb{R}) := \{v \in H^1(\mathbb{R}) : v(x) = v(-x), x \in \mathbb{R}\}$. We denote by X_{odd} the set of odd functions.

We denote by $\mathcal{G}$ the star graph constituted by N half-lines attached to a common vertex $\nu = 0$. On the graph we define the spaces

$$L^p(\mathcal{G}) = \bigoplus_{j=1}^{N} L^p(\mathbb{R}_+),\ p > 1, \quad H^1(\mathcal{G}) = \bigoplus_{j=1}^{N} H^1(\mathbb{R}_+), \quad H^2(\mathcal{G}) = \bigoplus_{j=1}^{N} H^2(\mathbb{R}_+).$$

Depending on the context we will use the following notations for different objects:

– by $||\cdot||$ we denote the norm in $L^2(\mathbb{R})$ or in $L^2(\mathcal{G})$ (accordingly $(\cdot,\cdot)$ denotes the scalar product in $L^2(\mathbb{R})$ or in $L^2(\mathcal{G})$);

– by $||\cdot||_p$ we denote the norm in $L^p(\mathbb{R})$ or in $L^p(\mathcal{G})$.

We also denote by $\mathcal{E}(\mathcal{G})$ the space $\mathcal{E}(\mathcal{G}) = \{\mathbf{V} = (v_j)_{j=1}^N \in H^1(\mathcal{G}) : v_1(0) = ... = v_N(0)\}$. For N even we define the space

$H^1_{\text{ev}}(\mathcal{G}) = \{\mathbf{V} = (v_j)_{j=1}^N \in H^1(\mathcal{G}) : v_1(x) = ... = v_{N/2}(x),\ v_{N/2+1}(x) = ... = v_N(x),\ x > 0\}$.

Let A be a densely defined symmetric operator in the Hilbert space $\mathcal{H}$. The domain of A is denoted by $\text{dom}(A)$. The deficiency numbers of A are denoted by $n_\pm(A) := \dim\ker(A^* \mp iI)$. The number of negative eigenvalues (counting multiplicities) is denoted by $n(A)$.

2 Preliminaries

2.1 NLS with point interactions.

Point interaction for (1.1) at $x = 0$ is formally defined by

$$\begin{pmatrix} u(t,0+) \\ \partial_x u(t,0+) \end{pmatrix} = \tau \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} u(t,0-) \\ \partial_x u(t,0-) \end{pmatrix}, \quad t \in \mathbb{R}, \tag{2.1}$$

in which

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{R}), \quad \text{and} \quad \tau \in S^1 \subset \mathbb{C}. \tag{2.2}$$

The equation (2.1) captures the interaction of the “field” u with the impurity (see [19]). The parameters in (2.2) label the self-adjoint extensions of the closable symmetric operator $H_0 = -\frac{d^2}{dx^2}$ defined, for instance, on the space $C_0^\infty(\mathbb{R}\setminus\{0\})$ of smooth functions with compact support separated from the origin $x = 0$. In fact, due to the theory of extensions of symmetric operators on Hilbert spaces, there exists a 4-parameter family of self-adjoint extensions which describes all one-point interactions of H_0. Such family can be equivalently described by means of boundary conditions at the origin

$$\begin{pmatrix} \psi(0+) \\ \psi'(0+) \end{pmatrix} = \tau \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \psi(0-) \\ \psi'(0-) \end{pmatrix}, \tag{2.3}$$

with a, b, c, d and τ satisfying the conditions in (2.2) (see [8, Theorem 3.2.3] or formula (K.1.2) from [7, Appendix K]).

We are interested in two specific choices of the parameters in (2.2), which are relevant in physical applications (see [3,19]). The first choice $\tau = a = d = 1$, $b = 0$, $c = -\gamma, \gamma \in \mathbb{R}\setminus\{0\}$,

$\mu = \mp 1$ corresponds to the δ-interaction of strength $-\gamma$ which gives rise to the following NLS-δ model

$$i\partial_t u - H_\gamma^\delta u \pm |u|^{p-1}u = 0, \tag{2.4}$$

where H_γ^δ is the self-adjoint operator on $L^2(\mathbb{R})$ defined by

$$\begin{aligned} &(H_\gamma^\delta v)(x) = -v''(x), \quad x \neq 0, \\ D_\gamma := &\operatorname{dom}(H_\gamma^\delta) = \big\{v \in H^1(\mathbb{R}) \cap H^2(\mathbb{R}\setminus\{0\}) : v'(0+) - v'(0-) = -\gamma v(0)\big\}. \end{aligned} \tag{2.5}$$

The operator H_γ^δ is formally defined by the expression $l_\gamma^\delta = -\frac{d^2}{dx^2} - \gamma\delta(x)$, where $\delta(x)$ is the Dirac delta distribution.

Equation (2.4) has been considered in various physical models with a point defect, for instance, in nonlinear optics and Bose-Einstein condensates. Indeed, the Dirac distribution models an impurity or defect localized at the origin. Moreover, NLS-δ equation can be viewed as a prototype model for the interaction of a wide soliton with a highly localized potential. In nonlinear optics it models a soliton propagating in a medium with a point defect, or interaction of a wide soliton with a much narrower one in a bimodal fiber (see [6, 13, 18, 25, 28, 29, 32, 33, 37, 38] and references therein).

The second choice of parameters $\tau = a = d = 1$, $c = 0$, $b = -\beta, \beta \in \mathbb{R}\setminus\{0\}$, $\mu = -1$ corresponds to the case of so-called δ'-interaction of strength $-\beta$. It gives rise to the following model (NLS-δ' henceforth)

$$i\partial_t u - H_\beta^{\delta'} u + |u|^{p-1}u = 0, \tag{2.6}$$

in which $H_\beta^{\delta'}$ is the self-adjoint operator on $L^2(\mathbb{R})$ defined by

$$\begin{aligned} &(H_\beta^{\delta'} v)(x) = -v''(x), \quad x \neq 0, \\ D_\beta := &\operatorname{dom}(H_\beta^{\delta'}) = \{v \in H^2(\mathbb{R}\setminus\{0\}) : v(0+) - v(0-) = -\beta v'(0),\ v'(0+) = v'(0-)\}. \end{aligned}$$

Recall that $H_\beta^{\delta'}$ is formally defined by the expression $l_\beta^{\delta'} = -\frac{d^2}{dx^2} - \beta < \cdot, \delta' > \delta'(x)$.

As far as we know, NLS-δ' model is less studied. In particular, the problem of the well-posedness of the dynamics is treated in [5] for the whole family of point interactions and cubic nonlinearity. Moreover, in [3, 4] the authors study variational properties and orbital stability of the ground states of the NLS-δ' equation.

The second part of our paper is devoted to investigation of the following nonlinear Schrödinger equation on the star graph $\mathcal{G}$

$$i\partial_t \mathbf{U}(t,x) + \partial_x^2 \mathbf{U}(t,x) + |\mathbf{U}(t,x)|^{p-1}\mathbf{U}(t,x) = 0, \tag{2.7}$$

where $\mathbf{U}(t,x) = (u_j(t,x))_{j=1}^N : \mathbb{R}\times\mathbb{R} \to \mathbb{C}^N$ and $p > 1$. Moreover, $\mathbf{U}(t,x)$ satisfies the following impurity conditions at $x = 0$

$$(U - I)\mathbf{U}(t,0) + i(U + I)\mathbf{U}'(t,0) = 0, \tag{2.8}$$

where $\mathbf{U}(t,0) = (u_j(t,0))_{j=1}^N$, $\mathbf{U}'(t,0) = (u_j'(t,0))_{j=1}^N$ and U is an arbitrary unitary $N \times N$ matrix and I is the $N \times N$ identity matrix . The conditions (2.8) at $x = 0$ define the N^2-parameter family of self-adjoint extensions of the closable symmetric operator ([16, Chapter 17])

$$\mathbf{H}_0 = \bigoplus_{j=1}^N \frac{-d^2}{dx^2}, \quad \operatorname{dom}(\mathbf{H}_0) = \bigoplus_{j=1}^N C_0^\infty(\mathbb{R}_+).$$

Equation (2.7) models propagation through junctions in networks (see [13,31]). The analysis of the behavior of NLS on networks is not yet fully developed, but it is currently growing (see [1, 2] and references therein). The linear Schrödinger equation on graphs has been a very developed subject due to its applications in quantum chemistry, nanotechnologies, and mesoscopic physics (see [16] for details and references).

Similarly to the previous scalar case we consider two choices of matrix U which correspond to so-called δ- and δ'- interactions on the star graph $\mathcal{G}$. More precisely, the matrix

$$U = \frac{2}{N + i\alpha}\mathcal{I} - I, \qquad \alpha \in \mathbb{R} \setminus \{0\},$$

where $\mathcal{I}$ is the $N \times N$ matrix whose all entries equal one, induces the following nonlinear Schrödinger equation with δ-interaction on the graph $\mathcal{G}$

$$i\partial_t \mathbf{U} - \mathbf{H}^\delta_\alpha \mathbf{U} + |\mathbf{U}|^{p-1}\mathbf{U} = 0. \tag{2.9}$$

Here $\mathbf{H}^\delta_\alpha$ is the self-adjoint operator on $L^2(\mathcal{G})$ defined for $\mathbf{V} = (v_j)_{j=1}^N$ by

$$\begin{aligned} (\mathbf{H}^\delta_\alpha \mathbf{V})(x) &= \left(-v_j''(x)\right)_{j=1}^N, \quad x > 0, \\ D_\alpha := \operatorname{dom}(\mathbf{H}^\delta_\alpha) &= \left\{ \mathbf{V} \in H^2(\mathcal{G}) : v_1(0) = \ldots = v_N(0), \ \sum_{j=1}^N v_j'(0) = \alpha v_1(0) \right\}. \end{aligned} \tag{2.10}$$

Model (2.9) has been extensively studied in [1, 2]. In particular, the authors showed well-posedness of the corresponding Cauchy problem. Moreover, they investigated the existence and the particular form of standing waves, as well as their variational and stability properties (see Theorems 2.2 and 2.4 below).

The second model we are interested in corresponds to

$$U = I - \frac{2}{N - i\lambda}\mathcal{I}, \qquad \lambda \in \mathbb{R} \setminus \{0\},$$

which induces nonlinear Schrödinger equation with δ'-interaction on the graph $\mathcal{G}$

$$i\partial_t \mathbf{U} - \mathbf{H}^{\delta'}_\lambda \mathbf{U} + |\mathbf{U}|^{p-1}\mathbf{U} = 0. \tag{2.11}$$

$\mathbf{H}^{\delta'}_\lambda$ is the self-adjoint operator on $L^2(\mathcal{G})$ defined for $\mathbf{V} = (v_j)_{j=1}^N$ by

$$\begin{aligned} (\mathbf{H}^{\delta'}_\lambda \mathbf{V})(x) &= \left(-v_j''(x)\right)_{j=1}^N, \quad x > 0, \\ D_\lambda := \operatorname{dom}(\mathbf{H}^{\delta'}_\lambda) &= \left\{ \mathbf{V} \in H^2(\mathcal{G}) : v_1'(0) = \ldots = v_N'(0), \ \sum_{j=1}^N v_j(0) = \lambda v_1'(0) \right\}. \end{aligned} \tag{2.12}$$

To our knowledge such type of interaction has never been studied for NLS on the star graph. In this connection one of the principal aims of this paper is to establish some results about the existence and stability of standing wave solutions of (2.11).

2.2 Review of the results on orbital stability for NLS equation with point interactions.

Crucial role in the stability analysis is played by the symmetries of the NLS equation (1.1). The basic symmetry associated to the mentioned equation is phase-invariance (in particular, translation invariance does not hold due to the defect). Thus, it is reasonable to define orbital stability as follows (for equations (2.4) and (2.6)).

Definition 2.1. The standing wave $u(t,x) = e^{i\omega t}\varphi(x)$ is said to be *orbitally stable* in a Hilbert space X if for any $\varepsilon > 0$ there exists $\eta > 0$ with the following property: if $u_0 \in X$ satisfies $||u_0 - \varphi||_X < \eta$, then the solution $u(t)$ of (2.4) (resp. (2.6)) with $u(0) = u_0$ exists for any $t \geq 0$ and

$$\sup_{t\in[0,\infty)} \inf_{\theta\in\mathbb{R}} ||u(t) - e^{i\theta}\varphi||_X < \varepsilon.$$

Otherwise, the standing wave $u(t,x) = e^{i\omega t}\varphi(x)$ is said to be *orbitally unstable* in X.

For NLS-δ and NLS-δ' equations on the line the space X coincides with $H^1(\mathbb{R})$ and $H^1(\mathbb{R} \setminus \{0\})$ respectively. Note also that the above definition is analogous in the case of NLS equation on the star graph $\mathcal{G}$, and the corresponding space X coincides with $\mathcal{E}(\mathcal{G})$ for equation (2.9) and with $H^1(\mathcal{G})$ for equation (2.11).

The existence of standing wave solutions of the equation (2.4) requires that the profile $\varphi = \varphi_{\omega,\gamma} \in D_\gamma$ satisfies the semi-linear elliptic equation

$$H^\delta_\gamma\varphi + \omega\varphi - |\varphi|^{p-1}\varphi = 0. \tag{2.13}$$

Fukuizumi and Jeanjean in [23] (see also [25]) showed that (2.13) for $\omega > \frac{\gamma^2}{4}$ has a unique positive even solution modulo rotation (see Figure 1)

$$\varphi_{\omega,\gamma}(x) = \left[\frac{(p+1)\omega}{2}\operatorname{sech}^2\left(\frac{(p-1)\sqrt{\omega}}{2}|x| + \tanh^{-1}\left(\frac{\gamma}{2\sqrt{\omega}}\right)\right)\right]^{\frac{1}{p-1}}, \quad x \in \mathbb{R}. \tag{2.14}$$



Figure 1(a) : $\varphi_{\omega,\gamma}(x)$ for $\gamma > 0$



Figure 1(b): $\varphi_{\omega,\gamma}(x)$ for $\gamma < 0$

This solution is constructed from the known solution of (2.13) in the case $\gamma = 0$ on each side of the defect pasted together at $x = 0$ to satisfy the condition of continuity and the jump condition $\varphi'(0+) - \varphi'(0-) = -\gamma\varphi(0)$ determined by (2.5). Note also that there is no nontrivial solution of (2.13) for $\omega \leq \frac{\gamma^2}{4}$ (see, for instance, [4, Appendix 8] for detailed explanation of this fact).

For the sake of completeness, we recall the main results on the stability of soliton solutions of (2.4). For $\gamma = 0$ orbital stability has been extensively studied in [14, 20, 21, 39]. Namely, $e^{i\omega t}\varphi_{\omega,0}$ is stable in $H^1(\mathbb{R})$ for any $\omega > 0$ and $1 < p < 5$ (see [20]) and unstable in $H^1(\mathbb{R})$ for any $\omega > 0$ and $p \geq 5$ (see [14] for $p > 5$ and [39] for $p = 5$).

The case $\gamma > 0$ was studied in [24]. In particular, authors showed that the standing wave $e^{i\omega t}\varphi_{\omega,\gamma}$ is stable in $H^1(\mathbb{R})$ for any $\omega > \frac{\gamma^2}{4}$ and $1 < p \leq 5$, and if $p > 5$, there exists a critical ω^* such that $e^{i\omega t}\varphi_{\omega,\gamma}$ is stable in $H^1(\mathbb{R})$ for any $\omega \in \left(\frac{\gamma^2}{4}, \omega^*\right)$ and unstable in $H^1(\mathbb{R})$ for any $\omega > \omega^*$. In the case $\gamma < 0$, the standing wave $e^{i\omega t}\varphi_{\omega,\gamma}$ is unstable "almost for sure" in $H^1(\mathbb{R})$ for any $p > 1$ (see [22, 23, 35]).

As it was mentioned above, the literature on stability of standing waves for NLS-δ' equations is very limited. As far as we know there are only two papers [3, 4] which establish extensive investigation on the topic. In [4] the case of $1 < p < 5$ and $\beta > 0$ was considered. Meanwhile, in [3] the authors investigated stability of standing waves for NLS-δ' equation in more general setting (in particular, for $p > 1$ and $\beta > 0$).

As above the existence of standing wave solutions $u(t,x) = e^{i\omega t}\varphi_{\omega,\beta}(x)$ of equation (2.6) requires that the profile $\varphi = \varphi_{\omega,\beta} \in D_\beta$ satisfies the semi-linear elliptic equation

$$H_\beta^{\delta'}\varphi + \omega\varphi - |\varphi|^{p-1}\varphi = 0. \tag{2.15}$$

It was shown in [3] that for $\beta > 0$ equation (2.15) has two types of solutions (odd and asymmetric, see Figure 2)

$$\varphi_{\omega,\beta}^{odd}(x) = \operatorname{sign}(x)\left[\frac{(p+1)\omega}{2}\operatorname{sech}^2\left(\frac{(p-1)\sqrt{\omega}}{2}(|x|+y)\right)\right]^{\frac{1}{p-1}}, \quad x \neq 0; \quad \tfrac{4}{\beta^2} < \omega, \tag{2.16}$$

$$\varphi_{\omega,\beta}^{as}(x) = \begin{cases} \left[\frac{(p+1)\omega}{2}\operatorname{sech}^2\left(\frac{(p-1)\sqrt{\omega}}{2}(x+y_1)\right)\right]^{\frac{1}{p-1}}, & x > 0; \\ -\left[\frac{(p+1)\omega}{2}\operatorname{sech}^2\left(\frac{(p-1)\sqrt{\omega}}{2}(x-y_2)\right)\right]^{\frac{1}{p-1}}, & x < 0, \end{cases}, \quad \omega > \frac{4}{\beta^2}\frac{p+1}{p-1},$$

where $y,\ y_1$ and y_2 are positive constants depending on β, p, ω (see [3, Theorem 5.3]).



Figure 2(a): $\varphi_{\omega,\beta}^{odd}$ for $\beta > 0$

Figure 2(b): $\varphi_{\omega,\beta}^{as}$ for $\beta > 0$

In [3, 4] were established the following stability results:

- standing wave $e^{i\omega t}\varphi^{odd}_{\omega,\beta}$ is stable for $p>1, \omega \in \left(\frac{4}{\beta^2}, \frac{4}{\beta^2}\frac{p+1}{p-1}\right)$ and unstable for $p>1$, $\omega > \frac{4}{\beta^2}\frac{p+1}{p-1}$;
- standing wave $e^{i\omega t}\varphi^{ass}_{\omega,\beta}$ is stable for $1<p\le 5,\ \omega > \frac{4}{\beta^2}\frac{p+1}{p-1}$, and $p>5,\ \omega \in \left(\frac{4}{\beta^2}\frac{p+1}{p-1}, \omega_1\right)$, meanwhile $e^{i\omega t}\varphi^{ass}_{\omega,\beta}$ is unstable for $p>5,\ \omega>\omega_2>\omega_1$.

The solutions in the case $\beta<0$ can be defined similarly (see Figure 3). In this case the constants y, y_1, y_2 have to be negative.



Figure 3(a): $\varphi^{odd}_{\omega,\beta}$ for $\beta<0$



Figure 3(b): $\varphi^{ass}_{\omega,\beta}$ for $\beta<0$

In the second part of the paper we study the orbital stability of the standing wave solutions

$$\mathbf{U}(t,x) = e^{i\omega t}\mathbf{\Phi}(x) = \left(e^{i\omega t}\varphi_j(x)\right)_{j=1}^N$$

for the NLS-δ and NLS-δ' equation on the star graph which appears to be more complicated problem. It is easily seen that amplitude $\mathbf{\Phi}\in D_\alpha$ of the standing wave of equation (2.9) satisfies the following stationary equation

$$\mathbf{H}^\delta_\alpha\mathbf{\Phi} + \omega\mathbf{\Phi} - |\mathbf{\Phi}|^{p-1}\mathbf{\Phi} = 0. \tag{2.17}$$

In [2] the authors obtained the following description of all solutions to equation (2.17).

Theorem 2.2. *[2, Theorem 4] Let* $[\cdot]$ *denote the integer part of the real number. Let also* $\alpha<0$, *and* $\omega > \frac{\alpha^2}{(N-2k)^2}$, *for some* $k\in\left\{0,...,\left[\frac{N-1}{2}\right]\right\}$. *Then equation* (2.17) *has* $k+1$ *(up to permutations) vector solutions* $\mathbf{\Phi}^\alpha_m = (\varphi^\alpha_{m,j})_{j=1}^N$, $m=0,...,k$, *which are given by*

$$\varphi^\alpha_{m,j}(x) = \begin{cases} \left[\frac{(p+1)\omega}{2}\operatorname{sech}^2\left(\frac{(p-1)\sqrt{\omega}}{2}x - a_m\right)\right]^{\frac{1}{p-1}}, & j=1,..,m; \\ \left[\frac{(p+1)\omega}{2}\operatorname{sech}^2\left(\frac{(p-1)\sqrt{\omega}}{2}x + a_m\right)\right]^{\frac{1}{p-1}}, & j=m+1,...,N, \end{cases}$$

$$\textit{where}\ \ a_m = \tanh^{-1}\left(\frac{\alpha}{(2m-N)\sqrt{\omega}}\right).$$

Moreover, for $-N\sqrt{\omega}<\alpha<\alpha^*<0$, *the vector solution* $\mathbf{\Phi}^\alpha_0 = (\varphi^\alpha_{0,j})_{j=1}^N$,

$$\varphi^\alpha_{0,j}(x) = \left[\frac{(p+1)\omega}{2}\operatorname{sech}^2\left(\frac{(p-1)\sqrt{\omega}}{2}x + \tanh^{-1}\left(\frac{-\alpha}{N\sqrt{\omega}}\right)\right)\right]^{\frac{1}{p-1}} \tag{2.18}$$

is the ground state.

Remark 2.3. Note that vector $\mathbf{\Phi}_m^\alpha = (\varphi_{m,j}^\alpha)_{j=1}^N$ has *m bumps* and $N-m$ *tails*. It is easily seen that $\mathbf{\Phi}_0^\alpha$ is the *N-tail profile*. Moreover, the N-tail profile is the only symmetric (i.e. invariant under permutations of the edges) solution of equation (2.17). In the case $N=5$ we have three types of profiles: a *5-tail profile*, a *4-tail/1-bump profile* and a *3-tail/2-bump profile*. They are demonstrated on Figure 4 (from the left to the right).



Figure 4

The parameter α^* in Theorem 2.2 originates from the variational problem associated with equation (2.17) and guarantees constrained minimality of the action functional

$$S_\omega^\alpha(\mathbf{\Psi}) = \tfrac{1}{2}||\mathbf{\Psi}'||^2 + \tfrac{\omega}{2}||\mathbf{\Psi}||^2 - \tfrac{1}{p+1}||\mathbf{\Psi}||_{p+1}^{p+1} + \tfrac{\alpha}{2}|\psi_1(0)|^2, \quad \mathbf{\Psi} = (\psi_j)_{j=1}^N \in \mathcal{E}(\mathcal{G}). \tag{2.19}$$

Namely, the vector solution $\mathbf{\Phi}_0^\alpha$ is the ground state in the sense of minimality of $S_\omega^\alpha(\mathbf{\Psi})$ at $\mathbf{\Phi}_0^\alpha$ with the constraint given by the Nehari manifold

$$\mathcal{N} = \{\mathbf{\Psi} \in \mathcal{E}(\mathcal{G}) : ||\mathbf{\Psi}'||^2 + \omega||\mathbf{\Psi}||^2 - ||\mathbf{\Psi}||_{p+1}^{p+1} + \alpha|\psi_1(0)|^2 = 0\}.$$

In [2] the following orbital stability result has been shown.

Theorem 2.4. *[2, Theorem 2] Let $1 < p \le 5$, $\alpha < \alpha^* < 0$, and $\omega > \frac{\alpha^2}{N^2}$. Then the ground state $e^{i\omega t}\mathbf{\Phi}_0^\alpha$ is orbitally stable in $\mathcal{E}(\mathcal{G})$.*

Stronger version of the above theorem was proved in [1, Theorem 1]. In particular, the authors proved orbital stability of $e^{i\omega t}\mathbf{\Phi}_0^\alpha$ for $\alpha < 0$ without restriction $\alpha < \alpha^* < 0$. The proof is based on a new original technique that allows one to restrict the analysis to functions made of pieces of soliton, reducing the problem to a finite dimensional one and avoiding to use direct methods of calculus of variations and linearization procedures (which is standard for Grillakis, Shatah and Strauss approach). Note also that, analogously to the scalar case, the authors in [2] showed that for $p > 5$ there exists ω^* such that $e^{i\omega t}\mathbf{\Phi}_0^\alpha$ is stable in $\mathcal{E}(\mathcal{G})$ for any $\omega \in \left(\frac{\alpha^2}{N^2}, \omega^*\right)$ and unstable for any $\omega > \omega^*$.

To our knowledge, the problem of orbital stability of standing waves $\mathbf{U}(t,x) = e^{i\omega t}\mathbf{\Phi}(x)$ has never been considered for NLS-δ' equation (2.11) on the star graph. In the present paper we study the orbital stability of the standing waves $\mathbf{U}(t,x) = e^{i\omega t}\mathbf{\Phi}_0^\lambda$ with tail-like profile $\mathbf{\Phi}_0^\lambda = (\varphi_0^\lambda)_{j=1}^N$, where

$$\varphi_0^\lambda(x) = \left[\frac{(p+1)\omega}{2}\operatorname{sech}^2\left(\frac{(p-1)\sqrt{\omega}}{2}x + \tanh^{-1}\left(\frac{-N}{\lambda\sqrt{\omega}}\right)\right)\right]^{\frac{1}{p-1}}, \quad \omega > \tfrac{N^2}{\lambda^2}, \quad \lambda < 0. \tag{2.20}$$

In Section 4 we prove a quite new result on stability of $e^{i\omega t}\mathbf{\Phi}_0^\lambda$ (see Theorem 1.1). This result in the case $N=2$ coincides with the known results for standing waves $e^{i\omega t}\varphi_{\omega,\beta}^{odd}(x)$ of NLS-δ' equation on the line.

Finalizing we note that investigation of the stability of the standing waves of NLS-δ and NLS-δ' equation on the star graph with bumps in profile $\mathbf{\Phi}(x)$ appears to be a complicated problem. The last one will be a topic of our future work.

3 Orbital stability of standing wave solutions for NLS-δ and NLS-δ' equation on the line

3.1 NLS-δ equation

Consider NLS-δ equation with attractive nonlinearity

$$i\partial_t u - H_\gamma^\delta u + |u|^{p-1}u = 0.$$

As it was mentioned above, one of the main approaches in stability analysis is the abstract theory by Grillakis, Shatah and Strauss [26, 27]. To formulate the stability criterium for the NLS-δ equation in the framework of this theory we define the following two self-adjoint linear operators

$$L_{1,\omega}^\gamma = -\frac{d^2}{dx^2} + \omega - p\varphi_{\omega,\gamma}^{p-1}, \quad L_{2,\omega}^\gamma = -\frac{d^2}{dx^2} + \omega - \varphi_{\omega,\gamma}^{p-1}, \quad \mathrm{dom}(L_{j,\omega}^\gamma) = D_\gamma,\ j\in\{1,2\}. \tag{3.1}$$

The operators $L_{1,\omega}^\gamma$ and $L_{1,\omega}^\gamma$ are associated with the key functional

$$S_\omega^\gamma(\psi) = \tfrac{1}{2}||\psi'||^2 + \tfrac{\omega}{2}||\psi||^2 - \tfrac{1}{p+1}||\psi||_{p+1}^{p+1} - \tfrac{\gamma}{2}|\psi(0)|^2$$

in the following sense

$$(S_\omega^\gamma)''(\varphi_{\omega,\gamma})(u,v) = (L_{1,\omega}^\gamma u_1, v_1) + (L_{2,\omega}^\gamma u_2, v_2),$$

where $u = u_1 + iu_2$ and $v = v_1 + iv_2$. The functions u_j, v_j, $j\in\{1,2\}$ are assumed to be real valued. Note also that $\varphi_{\omega,\gamma}$ is a critical point of S_ω^γ.

Since initial value problem associated to the NLS-δ equation is (at least) locally well-posed in $H^1(\mathbb{R})$ (see [21, Theorem 4.6.1]) and standing wave solution exists (that is, *Assumptions 1, 2* in [26] are satisfied), stability of the standing wave $e^{i\omega t}\varphi_{\omega,\gamma}(x)$ might be established by the following theorem (see [26, Theorem 2]).

Theorem 3.1. *Let $\omega > \frac{\gamma^2}{4}$. Suppose that the following spectral conditions hold.*

(*i*) $\ker(L_{1,\omega}^\gamma) = \{0\}$.

(*ii*) $\ker(L_{2,\omega}^\gamma) = \mathrm{span}\{\varphi_{\omega,\gamma}\}$.

(*iii*) $n(L_{1,\omega}^\gamma) = 1$.

(*iv*) *The rest of the spectrum of the operators $L_{1,\omega}^\gamma$ and $L_{2,\omega}^\gamma$ is positive and bounded away from zero.*

Then the standing wave $e^{i\omega t}\varphi_{\omega,\gamma}(x)$ *is stable in* $H^1(\mathbb{R})$ *if and only if* $\partial_\omega||\varphi_{\omega,\gamma}||^2$ *is positive.*

Making use of the explicit form (2.14) for $\varphi_{\omega,\gamma}$, the sign of $\partial_\omega||\varphi_{\omega,\gamma}||^2$ was computed in [23, 24]. By variational methods, it was shown in [23] that $n(L^\gamma_{1,\omega}) = 1$ in $H^1_{\text{rad}}(\mathbb{R})$, for arbitrary γ. Moreover, by using analytic perturbation theory and continuation argument, it was shown in [22] that $n(L^\gamma_{1,\omega}) = 1$ in $H^1(\mathbb{R})$ for any $\gamma > 0$, as well as $n(L^\gamma_{1,\omega}) = 2$ for $\gamma < 0$.

The equality $\ker(L^\gamma_{2,\omega}) = \operatorname{span}\{\varphi_{\omega,\gamma}\}$ was immediately deduced from (2.13) and simplicity of zero eigenvalue (see, for instance, [15, Chapter 2.3]). From the positivity of the profile $\varphi_{\omega,\gamma}$ it follows that $n(L^\gamma_{2,\omega}) = 0$ (see Lemma 3.3 below). The equality $\ker(L^\gamma_{1,\omega}) = \{0\}$ for any $\gamma \neq 0$ follows from the fact that $v(x) = \begin{cases} \mu\varphi'_{\omega,\gamma}, & x > 0 \\ \nu\varphi'_{\omega,\gamma}, & x < 0 \end{cases}$ does not belong to the domain D_γ as $\nu \neq 0, \mu \neq 0$. Positivity of the rest of the spectrum of the operators $L^\gamma_{1,\omega}$ and $L^\gamma_{2,\omega}$ is implied by the Weyl theorem on essential spectrum [36, Theorem XIII.14]. Thus, stability in the cases $H^1_{\text{rad}}(\mathbb{R})$, $\gamma \neq 0$, and $H^1(\mathbb{R})$, $\gamma > 0$, depends on the sign of $\partial_\omega||\varphi_{\omega,\gamma}||^2$.

Below we demonstrate how extension theory simplifies investigation of the spectrum of $L^\gamma_{1,\omega}$. In particular, we show the following key result.

Proposition 3.2. *Let* $L^\gamma_{1,\omega}$ *be defined in* (3.1) *and* $\omega > \frac{\gamma^2}{4}$. *Then* $n(L^\gamma_{1,\omega}) = 1$ *for any* $\gamma > 0$.

We establish two novel proofs of Proposition 3.2. The first one is based on a generalization of the classical Sturm oscillation theorem to the case of point interaction (see [9, 15]). We present its proof for completeness.

Lemma 3.3. *Let* $V(x)$ *be real-valued continuous function on* $\mathbb{R}$ *and* $\lim\limits_{|x|\to\infty} V(x) = c$. *Let also* $\varphi_1, \varphi_2 \in L^2(\mathbb{R})$ *be eigenfunctions of the operator*

$$L_V = -\frac{d^2}{dx^2} + V(x), \quad \operatorname{dom}(L_V) = D_\gamma,$$

corresponding to the eigenvalues $\lambda_1 < \lambda_2 < c$ *respectively. Suppose that* n_1 *and* n_2 *are the number of zeroes of* φ_1, φ_2 *respectively. Then* $n_2 > n_1$.

Proof. Suppose that $\varphi_1(a) = \varphi_1(b) = 0$ and $-\infty < a < 0 < b \leq \infty$, besides $\varphi_1(\infty) = 0$ is understood in the sense of limit. Let also $\varphi_1 > 0$ in (a, b). Then $\varphi_1'(a) > 0$ and $\varphi_1'(b) \leq 0$. The "equality" $\varphi_1'(b) = 0$ takes place only if $b = \infty$ since $\varphi_1 \in H^2(0, \infty)$. Suppose that φ_2 has no zeros in (a, b) and $\varphi_2 > 0$ in (a, b). Using the fact that φ_1, φ_2 are eigenfunctions of L_V, we arrive at

$$\begin{aligned}
0 &= \int_a^b (\varphi_1\varphi_2'' - \varphi_1''\varphi_2)dx + \int_a^b (\lambda_2 - \lambda_1)\varphi_1\varphi_2 dx \\
&= \int_a^{-0} \frac{d}{dx}(\varphi_1\varphi_2' - \varphi_1'\varphi_2)dx + \int_{+0}^b \frac{d}{dx}(\varphi_1\varphi_2' - \varphi_1'\varphi_2)dx + \int_a^b (\lambda_2 - \lambda_1)\varphi_1\varphi_2 dx \qquad (3.2) \\
&= [\varphi_1\varphi_2' - \varphi_1'\varphi_2]_a^b + [\varphi_1'\varphi_2 - \varphi_1\varphi_2']_{-0}^{+0} + \int_a^b (\lambda_2 - \lambda_1)\varphi_1\varphi_2 dx.
\end{aligned}$$

Since $\varphi_1, \varphi_2 \in D_\gamma$, we get $[\varphi_1'\varphi_2 - \varphi_1\varphi_2']_{-0}^{+0} = 0$. Therefore, from (3.2) and initial assumptions it easily follows that

$$0 > [\varphi_1\varphi_2' - \varphi_1'\varphi_2]_a^b = \varphi_1'(a)\varphi_2(a) - \varphi_1'(b)\varphi_2(b) > 0,$$

which is a contradiction. Thus, φ_2 has at least one zero in (a, b). Analogously, we can prove that there exists $\xi \in (-\infty, a]$ such that $\varphi_2(\xi) = 0$. Thereby, between two finite zeroes of φ_1 there exists a zero of φ_2 and between the last finite zero of φ_1 and ∞ (between the first finite zero of φ_1 and $-\infty$ respectively) there is at least one zero of φ_2. The proof is completed. □

The second proof of Proposition 3.2 uses extension theory of symmetric operators. For convenience of the reader and for the future references we formulate the following two results. The first one reads as follows (see [34, Chapter IV, §14]).

Proposition 3.4. *Let A be a densely defined lower semi-bounded symmetric operator (that is, $A \geq mI$) with finite deficiency indices $n_\pm(A) = k < \infty$ in the Hilbert space $\mathcal{H}$, and let $\widetilde{A}$ be a self-adjoint extension of A. Then the spectrum of $\widetilde{A}$ in $(-\infty, m)$ is discrete and consists of at most k eigenvalues counting multiplicities.*

Remark 3.5. (*i*) Proposition 3.4 holds for upper semi-bounded operator $A \leq MI$ and interval (M, ∞) respectively.

(*ii*) When $m = 0$, Proposition 3.4 provides an estimate for $n(\widetilde{A})$.

The second result is a corollary from the mini-max principle (see, for instance, [36, Chapter XIII]).

Proposition 3.6. *Let A be a self-adjoint operator in the Hilbert space $\mathcal{H}$ such that $\sigma_{\mathrm{ess}}(A) \subseteq [\omega, \infty)$, $\omega > 0$. Let there exist a finite-dimensional subspace $\mathcal{L}_-(A)$ such that the quadratic form of the operator A is negative on $\mathcal{L}_-(A) \setminus \{0\}$. Then $n(A) \geq \dim(\mathcal{L}_-(A))$.*

Remark 3.7. In connection with Lemma 3.3 and Proposition 3.6 it is worth mentioning spectral properties of the following self-adjoint operator corresponding to $\gamma = 0$ in (3.1)

$$L_{1,\omega}^0 = -\frac{d^2}{dx^2} + \omega - p\varphi_{\omega,0}^{p-1}(x), \qquad \mathrm{dom}(L_{1,\omega}^0) = H^2(\mathbb{R}).$$

Here $\varphi_{\omega,0}$ is the smooth soliton solution for the NLS equation (1.1) with $\mu = -1$ defined as

$$\varphi_{\omega,0}(x) = \left[\tfrac{(p+1)\omega}{2}\operatorname{sech}^2\left(\tfrac{(p-1)\sqrt{\omega}}{2}x\right)\right]^{\frac{1}{p-1}}. \tag{3.3}$$

The discrete spectrum of $L_{1,\omega}^0$ consists of two simple nonpositive distinct eigenvalues $\{\lambda_1, 0\}$. To see this one proves that the kernel of $L_{1,\omega}^0$ is spanned by $\varphi_{\omega,0}'$ which has only one zero. Hereafter one needs to apply Lemma 3.3 ($\gamma = 0$), Proposition 3.6, and the fact that $(L_{1,\omega}^0\varphi_{\omega,0}, \varphi_{\omega,0}) < 0$.

First proof of Proposition 3.2.

Initially we obtain from (2.13) that $(L_{1,\omega}^\gamma\varphi_{\omega,\gamma}, \varphi_{\omega,\gamma}) < 0$, and, due to Proposition 3.6, we have $n(L_{1,\omega}^\gamma) \geq 1$. To evaluate $n(L_{1,\omega}^\gamma)$ precisely consider the following self-adjoint operator

$$\widetilde{L}_{1,\omega}^\gamma = -\frac{d^2}{dx^2} + \omega - p\varphi_{\omega,0}^{p-1}, \qquad \mathrm{dom}(\widetilde{L}_{1,\omega}^\gamma) = D_\gamma,$$

where $\varphi_{\omega,0}$ is defined in (3.3). It is easily seen that $\varphi'_{\omega,0} \in \ker(\widetilde{L}^{\gamma}_{1,\omega})$. From Lemma 3.3 and the fact that $x = 0$ is the only zero of $\varphi'_{\omega,0}$ we have $n(\widetilde{L}^{\gamma}_{1,\omega}) \leq 1$. Since $\varphi_{\omega,0}(x) > \varphi_{\omega,\gamma}(x)$ for all $x \in \mathbb{R}$ and $\gamma > 0$ (**but not for** $\gamma < 0$ **!**), we get the following inequality

$$(L^{\gamma}_{1,\omega}v, v) \geq (\widetilde{L}^{\gamma}_{1,\omega}v, v), \quad \text{for all } \ v \in D_{\gamma}. \tag{3.4}$$

Combining inequality (3.4) with Proposition 3.6, we get

$$1 \leq n(L^{\gamma}_{1,\omega}) \leq n(\widetilde{L}^{\gamma}_{1,\omega}) \leq 1.$$

Thereby, in the case $\gamma > 0$ we get $n(L^{\gamma}_{1,\omega}) = 1$. □

The second of Proposition 3.2. Recall that $L^{\gamma}_{1,\omega}$ is the self-adjoint extension of the following symmetric operator

$$L_0 = -\frac{d^2}{dx^2} + \omega - p\varphi^{p-1}_{\omega,\gamma}, \quad \operatorname{dom}(L_0) = \left\{v \in H^2(\mathbb{R}) : \, v(0) = 0\right\}.$$

Moreover, it is known (see [7, Chapter I.3]) that

$$\operatorname{dom}(L_0^*) = H^1(\mathbb{R}) \cap H^2(\mathbb{R} \setminus \{0\}) = \operatorname{dom}(L_0) \dot{+} \operatorname{span}\{e^{i\sqrt{i}|x|}\} \dot{+} \operatorname{span}\{e^{i\sqrt{-i}|x|}\}, \quad \Im(\sqrt{\pm i}) > 0.$$

Next, it is easy to verify that for $\gamma > 0$ and $v \in H^2(\mathbb{R} \setminus \{0\})$ the following identity holds (see [3, Subsection 6.1])

$$-v'' + \omega v - p\varphi^{p-1}_{\omega,\gamma}v = \frac{-1}{\varphi'_{\omega,\gamma}}\frac{d}{dx}\left[(\varphi'_{\omega,\gamma})^2 \frac{d}{dx}\left(\frac{v}{\varphi'_{\omega,\gamma}}\right)\right], \qquad x \neq 0. \tag{3.5}$$

Then, using (3.5) and integrating by parts, we get

$$\begin{aligned}(L_0 v, v) = &\int\limits_{-\infty}^{0-} (\varphi'_{\omega,\gamma})^2 \left(\frac{d}{dx}\left(\frac{v}{\varphi'_{\omega,\gamma}}\right)\right)^2 dx \\ &+ \int\limits_{0+}^{\infty} (\varphi'_{\omega,\gamma})^2 \left(\frac{d}{dx}\left(\frac{v}{\varphi'_{\omega,\gamma}}\right)\right)^2 dx + \left[v'v - v^2 \frac{\varphi''_{\omega,\gamma}}{\varphi'_{\omega,\gamma}}\right]_{0-}^{0+}.\end{aligned} \tag{3.6}$$

The integral terms in (3.6) are non-negative and equal zero if and only if $v \equiv 0$. Due to the condition $v(0) = 0$, non-integral term vanishes, and we get $L_0 \geq 0$ on $\operatorname{dom}(L_0)$.

Due to the positivity of L_0, we conclude $n_-(L_0) = n_+(L_0)$ (see [34, Chapter IV,§14]) and, by Neumann formula (see [34, Chapter IV,§14]), deficiency indices of L_0 are given by

$$n_{\pm}(L_0) = \frac{\dim(\operatorname{dom}(L_0^*)/\operatorname{dom}(L_0))}{2} = 1.$$

Then, using Proposition 3.4 we get $n(L^{\gamma}_{1,\omega}) \leq 1$. This finishes the proof due to the inequality $(L^{\gamma}_{1,\omega}\varphi_{\omega,\gamma}, \varphi_{\omega,\gamma}) < 0$. □

Remark 3.8. The second proof of Proposition 3.2 cannot be applied for $\gamma < 0$ since in this case $\varphi'_{\omega,\gamma}$ has two zeroes.

NLS-δ equation with repulsive nonlinearity.

In [30] the authors considered stability of standing waves $u(t,x) = e^{i\omega t}\varphi_{\omega,\gamma}$ for the following nonlinear Schrödinger equation with repulsive nonlinearity on the line

$$i\partial_t u - H_\gamma^\delta u - |u|^{p-1}u = 0, \tag{3.7}$$

with the profile $\varphi = \varphi_{\omega,\gamma}$ satisfying the semi-linear elliptic equation

$$H_\gamma^\delta \varphi + \omega\varphi + |\varphi|^{p-1}\varphi = 0. \tag{3.8}$$

In [30] Kaminaga and Ohta using variational methods proved the following result.

Theorem 3.9. *[30, Theorem 1 and 2] Let $\gamma > 0$, then the following assertions hold.*

(i) *For $\omega = 0$ and $1 < p < 5$, the stationary problem* (3.8) *has unique positive solution (modulo rotations) $\varphi_{0,\gamma} \in H^1(\mathbb{R})$ given by*

$$\varphi_{0,\gamma} = \left[\frac{2(p+1)\gamma^2}{(4+(p-1)\gamma|x|)^2}\right]^{\frac{1}{p-1}}, \quad x \in \mathbb{R}.$$

Moreover, the standing wave solution $\varphi_{0,\gamma}$ is stable in $H^1(\mathbb{R})$.

(ii) *For $0 < \omega < \frac{\gamma^2}{4}$ and $p > 1$, the stationary problem* (3.8) *has unique positive solution (modulo rotations) $\varphi_{\omega,\gamma} \in H^1(\mathbb{R})$ given by*

$$\varphi_{\omega,\gamma} = \left[\frac{(p+1)\omega}{2}\sinh^{-2}\left(\frac{(p-1)\sqrt{\omega}}{2}|x| + \tanh^{-1}\left(\frac{2\sqrt{\omega}}{\gamma}\right)\right)\right]^{\frac{1}{p-1}}, \quad x \in \mathbb{R}. \tag{3.9}$$

Moreover, the standing wave solution $e^{i\omega t}\varphi_{\omega,\gamma}$ is stable in $H^1(\mathbb{R})$.

We recall that the authors also proved absence of nontrivial solution of (3.8) in the cases $\omega \notin \left[0, \frac{\gamma^2}{4}\right)$ and $\omega = 0,\ p \geq 5$.

The approach used in the first proof of Proposition 3.2 can be applied to give a short proof of the stability part of the above theorem. Let us show that the criterium of stability of Grillakis, Shatah and Strauss (see [26, Theorem 1]) might be easily applied. Indeed, for $0 < \omega < \frac{\gamma^2}{4}$ and $\gamma > 0$ we define the following two self-adjoint linear operators

$$L_{1,\omega}^\gamma = -\frac{d^2}{dx^2} + \omega + p\varphi_{\omega,\gamma}^{p-1}, \quad L_{2,\omega}^\gamma = -\frac{d^2}{dx^2} + \omega + \varphi_{\omega,\gamma}^{p-1}, \quad \mathrm{dom}(L_{j,\omega}^\gamma) = D_\gamma, \quad j \in \{1,2\}.$$

Here $\varphi_{\omega,\gamma}$ is defined by (3.9). Well-posedness of equation (3.7) in $H^1(\mathbb{R})$ was proved in [30, Propositon 1]. The equalities $\ker(L_{1,\omega}^\gamma) = \{0\}$, $\ker(L_{2,\omega}^\gamma) = \mathrm{span}\{\varphi_{\omega,\gamma}\}$, and positivity of the essential spectrum of the operators $L_{1,\omega}^\gamma, L_{2,\omega}^\gamma$ can be proved analogously to the previous case of attractive nonlinearity. Note also that the operators $L_{1,\omega}^\gamma$, $L_{2,\omega}^\gamma$ do not have negative eigenvalues. Indeed, since $\ker(L_{2,\omega}^\gamma) = \mathrm{span}\{\varphi_{\omega,\gamma}\}$ and $\varphi_{\omega,\gamma} > 0$, by Lemma 3.3, we get $n(L_{2,\omega}^\gamma) = 0$. Finally, equality $n(L_{1,\omega}^\gamma) = 0$ follows from the inequality $(L_{1,\omega}^\gamma v, v) > (L_{2,\omega}^\gamma v, v)$, $v \in D_\gamma \setminus \{0\}$. Since all the conditions of [26, Theorem 1] hold, we get the stability of the standing wave $e^{i\omega t}\varphi_{\omega,\gamma}(x)$ in $H^1(\mathbb{R})$.

The proof of stability of $\varphi_{0,\gamma}$ in $H^1(\mathbb{R})$ for $\omega = 0$ repeats the one in the case $0 < \omega < \frac{\gamma^2}{4}$.

3.2 NLS-δ' equation

As it was mentioned in the Preliminaries, orbital stability of standing wave solutions for NLS-δ' equation has been investigated in [3, 4]. In this Subsection we focus on the orbital stability of $e^{i\omega t}\varphi_{\omega,\beta}^{odd}$, where $\varphi_{\omega,\beta}^{odd}$ is defined by (2.16). In what follows we will use the notation $\varphi_{\omega,\beta}=\varphi_{\omega,\beta}^{odd}$.

Methods used in [4] are purely variational, while in [3] the authors combine variational approach and approach by Grillakis, Shatah and Strauss. Namely, they study spectral properties of the following two self-adjoint operators

$$\begin{aligned} &L_{1,\omega}^{\beta}=-\frac{d^2}{dx^2}+\omega-p|\varphi_{\omega,\beta}|^{p-1}, \quad L_{2,\omega}^{\beta}=-\frac{d^2}{dx^2}+\omega-|\varphi_{\omega,\beta}|^{p-1},\\ &\mathrm{dom}(L_{j,\omega}^{\beta})=D_\beta,\ \ j\in\{1,2\}. \end{aligned} \tag{3.10}$$

The operators $L_{1,\omega}^{\beta}$ and $L_{2,\omega}^{\beta}$ are associated with the action functional

$$S_\omega^\beta(\psi)=\tfrac12||\psi'||^2+\tfrac{\omega}{2}||\psi||^2-\tfrac{1}{p+1}||\psi||_{p+1}^{p+1}-\tfrac{1}{2\beta}|\psi(0+)-\psi(0-)|^2$$

in the sense of bilinear forms. Namely,

$$(S_\omega^\beta)''(\varphi_{\omega,\beta})(u,v)=(L_{1,\omega}^{\beta}u_1,v_1)+(L_{2,\omega}^{\beta}u_2,v_2),$$

where $u=u_1+iu_2$ and $v=v_1+iv_2$. The functions u_j, v_j, $j\in\{1,2\}$ are real valued.

The well-posedness for (2.6) in $H^1(\mathbb{R}\setminus\{0\})$ was established in [3, Proposition 3.3]. Note also that in the present case holds stability criterium analogous to Theorem 3.1 (the operators $L_{j,\omega}^{\gamma}$ have to be substituted by $L_{j,\omega}^{\beta}$). The spectral conditions $\ker(L_{1,\omega}^{\beta})=\{0\}$ and $\ker(L_{2,\omega}^{\beta})=\mathrm{span}\{\varphi_{\omega,\beta}\}$, for $\beta>0$, $\omega>\frac{4}{\beta^2}$, were established in [3]. The following proposition was proved in [3] by variational approach. We propose alternative proof in the framework of the extension theory

Proposition 3.10. *Let $\beta>0$, $L_{1,\omega}^{\beta}$ be defined in* (3.10)*, and $\omega>\frac{4}{\beta^2}$. Then*

(i) $n(L_{1,\omega}^{\beta})=1$ *for* $\omega\in\left(\frac{4}{\beta^2},\frac{4}{\beta^2}\frac{p+1}{p-1}\right]$.

(ii) $n(L_{1,\omega}^{\beta})=2$ *for* $\omega\in\left(\frac{4}{\beta^2}\frac{p+1}{p-1},\infty\right)$.

Proof. First, note that $L_{1,\omega}^{\beta}$ is the self-adjoint extension of the symmetric operator $L_{\min}$ defined by

$$L_{\min}=-\frac{d^2}{dx^2}+\omega-p|\varphi_{\omega,\beta}|^{p-1},\quad \mathrm{dom}(L_{\min})=\{v\in H^2(\mathbb{R}):v(0)=v'(0)=0\}.$$

To see that, one needs to put $\tau=a=d=1$, $b=0$ and $c=-\beta$ in (2.3). Since $\varphi_{\omega,\beta}\in L^\infty(\mathbb{R})$, we obtain

$$\mathrm{dom}(L_{\min}^*)=H^2(\mathbb{R}\setminus\{0\}).$$

Moreover, the operator $L_{\min}$ is non-negative for $\beta>0$. Indeed, it is easy to verify that for $\beta>0$ and $v\in H^2(\mathbb{R}\setminus\{0\})$ the following identity holds

$$-v''+\omega v-p|\varphi_{\omega,\beta}|^{p-1}v=\frac{-1}{\varphi_{\omega,\beta}'}\frac{d}{dx}\left[(\varphi_{\omega,\beta}')^2\frac{d}{dx}\left(\frac{v}{\varphi_{\omega,\beta}'}\right)\right],\quad x\neq 0, \tag{3.11}$$

where for $x \neq 0$

$$\varphi'_{\omega,\beta}(x) = \sqrt{\omega}\tanh\left(\tfrac{(p-1)\sqrt{\omega}}{2}(|x|+y)\right)\left[\tfrac{(p+1)\omega}{2}\operatorname{sech}^2\left(\tfrac{(p-1)\sqrt{\omega}}{2}(|x|+y)\right)\right]^{\frac{1}{p-1}}. \tag{3.12}$$

Here $y = \frac{1}{(p-1)\sqrt{\omega}}\ln\left(\frac{\beta\sqrt{\omega}+2}{\beta\sqrt{\omega}-2}\right) = \frac{2}{(p-1)\sqrt{\omega}}\tanh^{-1}\left(\frac{2}{\beta\sqrt{\omega}}\right)$ (see [3]). Using (3.11) and integrating by parts, we get

$$\begin{aligned}(L_{\min}v, v) = &\int\limits_{-\infty}^{0-}(\varphi'_{\omega,\beta})^2\left(\frac{d}{dx}\left(\frac{v}{\varphi'_{\omega,\beta}}\right)\right)^2 dx \\ &+\int\limits_{0+}^{\infty}(\varphi'_{\omega,\beta})^2\left(\frac{d}{dx}\left(\frac{v}{\varphi'_{\omega,\beta}}\right)\right)^2 dx + \left[v'v - v^2\frac{\varphi''_{\omega,\beta}}{\varphi'_{\omega,\beta}}\right]_{0-}^{0+}.\end{aligned} \tag{3.13}$$

The integral terms in (3.13) are non-negative and equal zero if and only if $v \equiv 0$. Due to the conditions $v(0) = v'(0) = 0$, non-integral term vanishes, and we get $L_{\min} \geq 0$.

Due to the positivity of $L_{\min}$ we conclude $n_-(L_{\min}) = n_+(L_{\min})$, and by Neumann formula deficiency indices of $L_{\min}$ are given by

$$n_\pm(L_{\min}) = \frac{\dim(\operatorname{dom}(L^*_{\min})/\operatorname{dom}(L_{\min}))}{2} = 2.$$

Indeed, (see, for instance, [7, Chapter 1.4])

$$H^2(\mathbb{R}\setminus\{0\}) = \operatorname{dom}(L_{\min})\dot{+}\operatorname{span}\{v^1_i, v^2_i\}\dot{+}\operatorname{span}\{v^1_{-i}, v^2_{-i}\},$$

where

$$v^1_{\pm i} = \begin{cases} e^{i\sqrt{\pm i}x}, & x > 0; \\ 0, & x < 0. \end{cases}, \qquad v^2_{\pm i} = \begin{cases} 0, & x > 0; \\ e^{-i\sqrt{\pm i}x}, & x < 0. \end{cases}, \quad \Im(\sqrt{\pm i}) > 0.$$

Since $L^\beta_{1,\omega}$ is the self-adjoint extension of the positive definite symmetric operator $L_{\min}$ and $n_\pm(L_{\min}) = 2$, by Proposition 3.4, $n(L^\beta_{1,\omega}) \leq 2$. Otherwise, we obtain from (2.15) that $(L^\beta_{1,\omega}\varphi_{\omega,\beta}, \varphi_{\omega,\beta}) < 0$, and, due to Proposition 3.6, we have $n(L^\beta_{1,\omega}) \geq 1$. Thus, we get $1 \leq n(L^\beta_{1,\omega}) \leq 2$.

(*i*) Note that $L^\beta_{1,\omega}$ is the self-adjoint extension of the following symmetric operator

$$L'_0 = -\frac{d^2}{dx^2} + \omega - p|\varphi_{\omega,\beta}|^{p-1}, \quad \operatorname{dom}(L'_0) = \left\{v \in H^2(\mathbb{R}) : v'(0) = 0\right\}.$$

Let us show that $L'_0 \geq 0$. Using (3.11) and integrating by parts, we get

$$\begin{aligned}(L'_0 v, v) = &\int\limits_{-\infty}^{0-}(\varphi'_{\omega,\beta})^2\left(\frac{d}{dx}\left(\frac{v}{\varphi'_{\omega,\beta}}\right)\right)^2 dx \\ &+\int\limits_{0+}^{\infty}(\varphi'_{\omega,\beta})^2\left(\frac{d}{dx}\left(\frac{v}{\varphi'_{\omega,\beta}}\right)\right)^2 dx + \left[v'v - v^2\frac{\varphi''_{\omega,\beta}}{\varphi'_{\omega,\beta}}\right]_{0-}^{0+}.\end{aligned} \tag{3.14}$$

The integral terms in (3.14) are non-negative. Let us focus on non-integral term. Due to the conditions $v'(0)=0, v(0+)=v(0-)$, and formula (3.12), we deduce

$$\left[v'v - v^2\frac{\varphi''_{\omega,\beta}}{\varphi'_{\omega,\beta}}\right]_{0-}^{0+} = -\left[v^2\frac{\varphi''_{\omega,\beta}}{\varphi'_{\omega,\beta}}\right]_{0-}^{0+} = v^2(0)\frac{\varphi''_{\omega,\beta}(0-)\varphi'_{\omega,\beta}(0+) - \varphi''_{\omega,\beta}(0+)\varphi'_{\omega,\beta}(0-)}{\varphi'_{\omega,\beta}(0+)\varphi'_{\omega,\beta}(0-)}$$
$$= v^2(0)\frac{\varphi''_{\omega,\beta}(0-) - \varphi''_{\omega,\beta}(0+)}{\varphi'_{\omega,\beta}(0-)} = -v^2(0)\frac{\beta\omega}{2}\left(p-1-(p+1)\frac{4}{\beta^2\omega}\right) \geq 0.$$

The last inequality follows from $\omega \leq \frac{4}{\beta^2}\frac{p+1}{p-1}$.

We have the following Neumann decomposition (see [7, Chapter I.4])

$$\operatorname{dom}((L'_0)^*) = \left\{v \in H^2(\mathbb{R}\setminus\{0\}) : \ v'(0+) = v'(0-)\right\} = \operatorname{dom}(L'_0) \dot{+} \operatorname{span}\{v_i\} \dot{+} \operatorname{span}\{v_{-i}\},$$

where

$$v_{\pm i} = \begin{cases} e^{i\sqrt{\pm i}x}, & x>0, \\ -e^{-i\sqrt{\pm i}x}, & x<0, \end{cases}, \quad \Im(\sqrt{\pm i}) > 0.$$

Due to the positivity of L'_0 and the above decomposition, we obtain

$$n_\pm(L'_0) = \frac{\dim(\operatorname{dom}((L'_0)^*)/\operatorname{dom}(L'_0))}{2} = 1.$$

By Proposition 3.4, we get $n(L^\beta_{1,\omega}) \leq 1$, and finally $n(L^\beta_{1,\omega}) = 1$.

(*ii*) The quadratic form of the operator $L^\beta_{1,\omega}$ is defined in $H^1(\mathbb{R}\setminus\{0\})$ as follows

$$F^\beta_{1,\omega}(u) = ||u'||^2 + \omega||u||^2 - p(|\varphi_{\omega,\beta}|^{p-1}u, u) - \tfrac{1}{\beta}|u(0+) - u(0-)|^2.$$

Noting that $\varphi'_{\omega,\beta}(0+) = \varphi'_{\omega,\beta}(0-)$ and integrating by parts, we get for $u = \varphi'_{\omega,\beta}$

$$F^\beta_{1,\omega}(\varphi'_{\omega,\beta}) = \int\limits_{\mathbb{R}_-}\varphi'_{\omega,\beta}\Big(-\varphi'''_{\omega,\beta} + \omega\varphi'_{\omega,\beta} - p|\varphi_{\omega,\beta}|^{p-1}\varphi'_{\omega,\beta}\Big)dx$$
$$+\int\limits_{\mathbb{R}_+}\varphi'_{\omega,\beta}\Big(-\varphi'''_{\omega,\beta} + \omega\varphi'_{\omega,\beta} - p|\varphi_{\omega,\beta}|^{p-1}\varphi'_{\omega,\beta}\Big)dx + \varphi'_{\omega,\beta}(0+)(\varphi''_{\omega,\beta}(0-) - \varphi''_{\omega,\beta}(0+))$$
$$= \varphi'_{\omega,\beta}(0+)(\varphi''_{\omega,\beta}(0-) - \varphi''_{\omega,\beta}(0+))$$
$$= -\tfrac{2}{\beta}\omega\left(\tfrac{(p+1)\omega}{2}\right)^{\frac{2}{p-1}}\left(1-\tfrac{4}{\beta^2\omega}\right)^{\frac{2}{p-1}}\left(p-1-(p+1)\tfrac{4}{\beta^2\omega}\right).$$

The last one expression is negative due to $\omega > \frac{4}{\beta^2}\frac{p+1}{p-1}$. Since $F^\beta_{1,\omega}(\varphi_{\omega,\beta}) = \left(L^\beta_{1,\omega}\varphi_{\omega,\beta}, \varphi_{\omega,\beta}\right) < 0$, and the functions $\varphi_{\omega,\beta}, \varphi'_{\omega,\beta}$ have different parity, we obtain for $r, s \in \mathbb{C}$

$$F^\beta_{1,\omega}(s\varphi_{\omega,\beta} + r\varphi'_{\omega,\beta}) = |s|^2F^\beta_{1,\omega}(\varphi_{\omega,\beta}) + |r|^2F^\beta_{1,\omega}(\varphi'_{\omega,\beta}) < 0.$$

Therefore, $F^\beta_{1,\omega}$ is negative on two-dimensional subspace $\mathcal{M} = \operatorname{span}\{\varphi_{\omega,\beta}, \varphi'_{\omega,\beta}\} \subset H^1(\mathbb{R}\setminus\{0\})$. Thus, Proposition 3.6 induces $n(L^\beta_{1,\omega}) \geq 2$, and consequently $n(L^\beta_{1,\omega}) = 2$.

□

In [3, Proposition 6.5] it was shown that $\partial_\omega ||\varphi_{\omega,\beta}||^2$ is positive for any $p > 1$ and $\omega \in \left(\frac{4}{\beta^2}, \frac{4}{\beta^2}\frac{p+1}{p-1}\right)$. Thus, due to Proposition 3.10, we conclude that $e^{i\omega t}\varphi_{\omega,\beta}$ is orbitally stable in this case.

Below we briefly discuss how to demonstrate instability of $e^{i\omega t}\varphi_{\omega,\beta}$ for $p > 1$ and $\omega > \frac{4}{\beta^2}\frac{p+1}{p-1}$ proved in [3, Theorem 6.11]. To do that we need the following key result.

Proposition 3.11. *Let* $\omega > \frac{4}{\beta^2}$, $\beta > 0$, *and operator* $\widetilde{L}^{\beta}_{1,\omega}$ *be defined as*

$$\widetilde{L}^{\beta}_{1,\omega} = -\frac{d^2}{dx^2} + \omega - p|\varphi_{\omega,\beta}|^{p-1}, \quad \mathrm{dom}(\widetilde{L}^{\beta}_{1,\omega}) = D_\beta \cap X_{\mathrm{odd}}.$$

Then $n(\widetilde{L}^{\beta}_{1,\omega}) = 1$.

Proof. It is obvious that $n(\widetilde{L}^{\beta}_{1,\omega}) \le n(L^{\beta}_{1,\omega}) = 1$. Since $\varphi_{\omega,\beta} \in \mathrm{dom}(\widetilde{L}^{\beta}_{1,\omega})$ and $(\widetilde{L}^{\beta}_{1,\omega}\varphi_{\omega,\beta}, \varphi_{\omega,\beta}) < 0$, by Proposition 3.6, we get $n(\widetilde{L}^{\beta}_{1,\omega}) = 1$. □

Well-posedness of the Cauchy problem in $H^1(\mathbb{R}) \cap X_{\mathrm{odd}}$ associated with (2.6) was shown in [3, Theorem 6.11]. Thus, by [27, Instability Theorem] we induce instability of $e^{i\omega t}\varphi_{\omega,\beta}$ for $p > 1$ and $\omega > \frac{4}{\beta^2}\frac{p+1}{p-1}$. Indeed, when $\partial_\omega||\varphi_{\omega,\beta}||^2 > 0$ then instability follows from Proposition 3.10(*ii*). In the case $\partial_\omega||\varphi_{\omega,\beta}||^2 < 0$ we can conclude by Proposition 3.11 instability of $e^{i\omega t}\varphi_{\omega,\beta}$ in $H^1(\mathbb{R}) \cap X_{\mathrm{odd}}$ which naturally induces instability in $H^1(\mathbb{R})$. If $\partial_\omega||\varphi_{\omega,\beta}||^2(\omega) = 0$, then instability in $H^1(\mathbb{R}) \cap X_{\mathrm{odd}}$ (and consequently in $H^1(\mathbb{R})$) follows from [35, Colollary 2] .

4 Orbital stability of standing wave solutions for NLS-δ and NLS-δ' equation on a star graph

4.1 NLS-δ equation on a star graph

In this Subsection we propose an alternative proof of the orbital stability of $e^{i\omega t}\mathbf{\Phi}^{\alpha}_0$ defined in (2.18) for $\alpha < 0$ (see Theorem 2.4 above and [1, Theorem 1]). Consider the following two self-adjoint matrix operators

$$\mathbf{L}^{\alpha}_{1,\omega} = \left(\left(-\frac{d^2}{dx^2} + \omega - p(\varphi^{\alpha}_0)^{p-1}\right)\delta_{k,j}\right), \quad \mathbf{L}^{\alpha}_{2,\omega} = \left(\left(-\frac{d^2}{dx^2} + \omega - (\varphi^{\alpha}_0)^{p-1}\right)\delta_{k,j}\right),$$
$$\mathrm{dom}(\mathbf{L}^{\alpha}_{1,\omega}) = \mathrm{dom}(\mathbf{L}^{\alpha}_{2,\omega}) = D_\alpha,$$

where $\delta_{k,j}$ is the Kronecker symbol, D_α is defined by (2.10), and $\varphi^{\alpha}_0 = \varphi^{\alpha}_{0,j}$ is defined in (2.18). Analogously to the scalar case the operators $\mathbf{L}^{\alpha}_{1,\omega}$ and $\mathbf{L}^{\alpha}_{2,\omega}$ are associated with the functional S^{α}_{ω} defined by (2.19), i.e.,

$$(S^{\alpha}_{\omega})''(\mathbf{\Phi}^{\alpha}_0)(\mathbf{U}, \mathbf{V}) = (\mathbf{L}^{\alpha}_{1,\omega}\mathbf{U}_1, \mathbf{V}_1) + (\mathbf{L}^{\alpha}_{2,\omega}\mathbf{U}_2, \mathbf{V}_2),$$

where $\mathbf{U} = \mathbf{U}_1 + i\mathbf{U}_2$ and $\mathbf{V} = \mathbf{V}_1 + i\mathbf{V}_2$. The vector functions $\mathbf{U}_j, \mathbf{V}_j$, $j \in \{1, 2\}$ are assumed to be real valued. Since *Assumptions 1, 2* in [26] hold (see [2, Proposition 2.1 and Theorem 4]), the following stability criterium holds.

Theorem 4.1. *Let $\alpha < 0$, $\omega > \frac{\alpha^2}{N^2}$, and the following spectral conditions hold.*

(*i*) $\ker(\mathbf{L}^{\alpha}_{1,\omega}) = \{\mathbf{0}\}$.

(*ii*) $\ker(\mathbf{L}^{\alpha}_{2,\omega}) = \operatorname{span}\{\mathbf{\Phi}^{\alpha}_0\}$.

(*iii*) $n(\mathbf{L}^{\alpha}_{1,\omega}) = 1$.

(*iv*) *The rest of the spectrum of the operators $\mathbf{L}^{\alpha}_{1,\omega}$ and $\mathbf{L}^{\alpha}_{2,\omega}$ is positive and bounded away from zero.*

Then the standing wave $e^{i\omega t}\mathbf{\Phi}^{\alpha}_0$ is stable in $\mathcal{E}(\mathcal{G})$ if and only if $\partial_\omega||\mathbf{\Phi}^{\alpha}_0||^2$ is positive.

The conditions (*i*), (*ii*), (*iv*) of Theorem 4.1 were established in [2]. The condition (*iii*) was proved for $-N\sqrt{\omega} < \alpha < \alpha^*$ using variational approach. We prove that $n(\mathbf{L}^{\alpha}_{1,\omega}) = 1$ in the framework of the extension theory for $\alpha < 0$ without restriction $\alpha < \alpha^*$.

Proposition 4.2. *Let $\omega > \frac{\alpha^2}{N^2}$ and $\alpha < 0$. Then $n(\mathbf{L}^{\alpha}_{1,\omega}) = 1$.*

Proof. In what follows we will use the notation $\boldsymbol{l}_\alpha = \Big(\Big(-\frac{d^2}{dx^2} + \omega - p(\varphi^{\alpha}_{0,j})^{p-1}\Big)\delta_{k,j}\Big)$. First, note that $\mathbf{L}^{\alpha}_{1,\omega}$ is the self-adjoint extension of the following symmetric operator

$$\mathbf{L}_0 = \boldsymbol{l}_\alpha, \quad \operatorname{dom}(\mathbf{L}_0) = \left\{\mathbf{V} \in H^2(\mathcal{G}) : v_1(0) = ... = v_N(0) = 0, \sum_{j=1}^{N} v_j'(0) = 0\right\}.$$

Below we show that the operator $\mathbf{L}_0$ is non-negative and $n_\pm(\mathbf{L}_0) = 1$. First, let us show that the adjoint operator of $\mathbf{L}_0$ is given by

$$\mathbf{L}_0^* = \boldsymbol{l}_\alpha, \quad \operatorname{dom}(\mathbf{L}_0^*) = \big\{\mathbf{V} \in H^2(\mathcal{G}) : v_1(0) = ... = v_N(0)\big\} =: D_0^*. \tag{4.1}$$

Using standard arguments one can prove that $\operatorname{dom}(\mathbf{L}_0^*) \subset H^2(\mathcal{G})$ and $\mathbf{L}_0^* = \boldsymbol{l}_\alpha$ (see [34, Chapter V,§17]). It is easily seen the inclusion $D_0^* \subseteq \operatorname{dom}(\mathbf{L}_0^*)$. Indeed, for any $\mathbf{U} = (u_j)_{j=1}^N \in D_0^*$ and $\mathbf{V} = (v_j)_{j=1}^N \in \operatorname{dom}(\mathbf{L}_0)$ denoting $\mathbf{U}^* = \boldsymbol{l}_\alpha(\mathbf{U}) \in L^2(\mathcal{G})$ we get

$$\begin{aligned}(\mathbf{L}_0\mathbf{V}, \mathbf{U}) &= (\boldsymbol{l}_\alpha(\mathbf{V}), \mathbf{U}) = (\mathbf{V}, \boldsymbol{l}_\alpha(\mathbf{U})) + \sum_{j=1}^{N}\left[-v_j'u_j + v_ju_j'\right]_0^\infty \\ &= (\mathbf{V}, \boldsymbol{l}_\alpha(\mathbf{U})) = (\mathbf{V}, \mathbf{U}^*),\end{aligned}$$

which, by definition of the adjoint operator, means that $\mathbf{U} \in \operatorname{dom}(\mathbf{L}_0^*)$ or $D_0^* \subseteq \operatorname{dom}(\mathbf{L}_0^*)$.

Let us show the inverse inclusion $D_0^* \supseteq \operatorname{dom}(\mathbf{L}_0^*)$. Take $\mathbf{U} \in \operatorname{dom}(\mathbf{L}_0^*)$, then for any $\mathbf{V} \in \operatorname{dom}(\mathbf{L}_0)$ we have

$$\begin{aligned}(\mathbf{L}_0\mathbf{V}, \mathbf{U}) &= (\boldsymbol{l}_\alpha(\mathbf{V}), \mathbf{U}) = (\mathbf{V}, \boldsymbol{l}_\alpha(\mathbf{U})) + \sum_{j=1}^{N}\left[-v_j'u_j + v_ju_j'\right]_0^\infty \\ &= (\mathbf{V}, \mathbf{L}_0^*\mathbf{U}) = (\mathbf{V}, \boldsymbol{l}_\alpha(\mathbf{U})).\end{aligned}$$

Thus, we arrive at the equality

$$\sum_{j=1}^{N}\left[-v_j'u_j+v_ju_j'\right]_0^{\infty}=\sum_{j=1}^{N}v_j'(0)u_j(0)=0 \tag{4.2}$$

for any $\mathbf{V}\in\operatorname{dom}(\mathbf{L}_0)$. Let $\mathbf{W}=(w_j)_{j=1}^N\in\operatorname{dom}(\mathbf{L}_0)$ such that $w_3'(0)=w_4'(0)=...=w_N'(0)=0$. Then for $\mathbf{U}\in\operatorname{dom}(\mathbf{L}_0^*)$ from (4.2) it follows that

$$\sum_{j=1}^{N}w_j'(0)u_j(0)=w_1'(0)u_1(0)+w_2'(0)u_2(0)=0. \tag{4.3}$$

Recalling that $\sum\limits_{j=1}^{N}w_j'(0)=w_1'(0)+w_2'(0)=0$ and assuming $w_2'(0)\neq 0$, we obtain from (4.3) the equality $u_1(0)=u_2(0)$. Repeating the similar arguments for $\mathbf{W}=(w_j)_{j=1}^N\in\operatorname{dom}(\mathbf{L}_0)$ such that $w_4'(0)=w_5'(0)=...=w_N'(0)\equiv 0$, we get $u_1(0)=u_2(0)=u_3(0)$ and so on. Finally taking $\mathbf{W}=(w_j)_{j=1}^N\in\operatorname{dom}(\mathbf{L}_0)$ such that $w_N'(0)=0$ we will arrive at $u_1(0)=u_2(0)=...=u_{N-1}(0)$ and consequently $u_1(0)=u_2(0)=...=u_N(0)$. Thus, $\mathbf{U}\in D_0^*$ or $D_0^*\supseteq\operatorname{dom}(\mathbf{L}_0^*)$, and (4.1) holds.

Let us show that the operator $\mathbf{L}_0$ is non-negative. First, note that every component of the vector $\mathbf{V}=(v_j)_{j=1}^N\in H^2(\mathcal{G})$ satisfies the following identity

$$-v_j''+\omega v_j-p(\varphi_{0,j}^{\alpha})^{p-1}v_j=\frac{-1}{(\varphi_{0,j}^{\alpha})'}\frac{d}{dx}\left[((\varphi_{0,j}^{\alpha})')^2\frac{d}{dx}\left(\frac{v_j}{(\varphi_{0,j}^{\alpha})'}\right)\right],\quad x\neq 0.$$

Using the above equality and integrating by parts, we get for $\mathbf{V}\in\operatorname{dom}(\mathbf{L}_0)$

$$\begin{aligned}(\mathbf{L}_0\mathbf{V},\mathbf{V})&=\sum_{j=1}^{N}\int_0^{\infty}((\varphi_{0,j}^{\alpha})')^2\left(\frac{d}{dx}\left(\frac{v_j}{(\varphi_{0,j}^{\alpha})'}\right)\right)^2dx+\sum_{j=1}^{N}\left[-v_j'v_j+v_j^2\frac{(\varphi_{0,j}^{\alpha})''}{(\varphi_{0,j}^{\alpha})'}\right]_0^{\infty}\\&=\sum_{j=1}^{N}\int_0^{\infty}((\varphi_{0,j}^{\alpha})')^2\left(\frac{d}{dx}\left(\frac{v_j}{(\varphi_{0,j}^{\alpha})'}\right)\right)^2dx\geq 0.\end{aligned}$$

Due to the Neumann formula we obtain the decomposition

$$\operatorname{dom}(\mathbf{L}_0^*)=\left\{\mathbf{V}\in H^2(\mathcal{G}):v_1(0)=...=v_N(0)\right\}=\operatorname{dom}(\mathbf{L}_0)\dotplus\operatorname{span}\{\mathbf{V}_i\}\dotplus\operatorname{span}\{\mathbf{V}_{-i}\},$$

where $\mathbf{V}_{\pm i}=\left(e^{i\sqrt{\pm i}x}\right)_{j=1}^N$. Indeed, since $\varphi_0^{\alpha}\in L^{\infty}(\mathbb{R}_+)$, we get $\operatorname{dom}(\mathbf{L}_0^*)=\operatorname{dom}(\mathbf{L}^*)$, where

$$\mathbf{L}=\left(\left(-\frac{d^2}{dx^2}\right)\delta_{k,j}\right),\quad \operatorname{dom}(\mathbf{L})=\operatorname{dom}(\mathbf{L}_0).$$

Therefore, deficiency indices are equal (by positivity of $\mathbf{L}_0$) and

$$n_{\pm}(\mathbf{L}_0)=\frac{\dim(\operatorname{dom}(\mathbf{L}_0^*)/\operatorname{dom}(\mathbf{L}_0))}{2}=1.$$

Due to Proposition 3.4, $n(\mathbf{L}_{1,\omega}^{\alpha})\leq 1$. Taking into account that $(\mathbf{L}_{1,\omega}^{\alpha}\mathbf{\Phi}_0^{\alpha},\mathbf{\Phi}_0^{\alpha})=-(p-1)||\mathbf{\Phi}_0^{\alpha}||_{p+1}^{p+1}<0$, we arrive at $n(\mathbf{L}_{1,\omega}^{\alpha})=1$. □

Theorem 2.4 now easily follows for $\alpha<0$ from Theorem 4.1 and positivity of $\partial_{\omega}||\mathbf{\Phi}_0^{\alpha}||^2$ proved in [2].

4.2 NLS-δ' equation on a star graph

As it was announced in the Introduction, in this Subsection we discuss a new problem. In particular, we study orbital stability of the standing wave $\mathbf{U}(t,x) = e^{i\omega t}\mathbf{\Phi}(x) = (e^{i\omega t}\varphi_j(x))_{j=1}^N$ of NLS-δ' equation (2.6) with the particular profile $\mathbf{\Phi} = \mathbf{\Phi}_0^\lambda = (\varphi_0^\lambda)_{j=1}^N$, where φ_0^λ is defined by (2.20) for $\lambda < 0$.

It is easily seen that amplitude $\mathbf{\Phi}_0^\lambda$ of the standing wave satisfies the stationary equation

$$\mathbf{H}_\lambda^{\delta'}\mathbf{\Phi} + \omega\mathbf{\Phi} - |\mathbf{\Phi}|^{p-1}\mathbf{\Phi} = 0.$$

In other words, standing waves exist or *Assumption 2* in [26] holds. As before we use the same approach by Grillakis, Shatah and Strauss. Due to this approach we need to prove well-posedness of the Cauchy problem associated with (2.11) (*Assumption 1* in [26]). The proof of this property can be easily established combining the ideas of proof of [3, Proposition 3.3] and [1, Proposition 2.1].

Consider two self-adjoint matrix operators

$$\mathbf{L}_{1,\omega}^\lambda = \left(\Big(-\frac{d^2}{dx^2} + \omega - p(\varphi_0^\lambda)^{p-1}\Big)\delta_{k,j}\right), \quad \mathbf{L}_{2,\omega}^\lambda = \left(\Big(-\frac{d^2}{dx^2} + \omega - (\varphi_0^\lambda)^{p-1}\Big)\delta_{k,j}\right),$$
$$\mathrm{dom}(\mathbf{L}_{1,\omega}^\lambda) = \mathrm{dom}(\mathbf{L}_{2,\omega}^\lambda) = D_\lambda,$$

where $\delta_{k,j}$ is the Kronecker symbol. These operators are associated in a standard way with the second derivative of the following action functional

$$S_\omega^\lambda(\mathbf{\Psi}) = \tfrac{1}{2}||\mathbf{\Psi}'||^2 - \tfrac{1}{p+1}||\mathbf{\Psi}||_{p+1}^{p+1} + \tfrac{1}{2\lambda}\left|\sum_{j=1}^N \psi_j(0)\right|^2 + \tfrac{\omega}{2}||\mathbf{\Psi}||^2,$$

where $\mathbf{\Psi} = (\psi_j)_{j=1}^N \in H^1(\mathcal{G})$. Namely,

$$(S_\omega^\lambda)''(\mathbf{\Phi}_0^\lambda)(\mathbf{U},\mathbf{V}) = (\mathbf{L}_{1,\omega}^\lambda\mathbf{U}_1, \mathbf{V}_1) + (\mathbf{L}_{2,\omega}^\lambda\mathbf{U}_2, \mathbf{V}_2),$$

where $\mathbf{U} = \mathbf{U}_1 + i\mathbf{U}_2$ and $\mathbf{V} = \mathbf{V}_1 + i\mathbf{V}_2$.

Since *Assumptions 1,2* in [27] are satisfied, the following theorem holds (see [27, Sability and Instability Theorem]).

Theorem 4.3. *Let $\lambda < 0, \omega > \frac{N^2}{\lambda^2}$, and the number $p(\omega_0)$ be defined as*

$$p(\omega_0) = \begin{cases} 1 & \text{if } \partial_\omega||\mathbf{\Phi}_0^\lambda||^2 > 0 \text{ at } \omega = \omega_0, \\ 0 & \text{if } \partial_\omega||\mathbf{\Phi}_0^\lambda||^2 < 0 \text{ at } \omega = \omega_0. \end{cases} \tag{4.4}$$

Let also the following spectral conditions hold.

(*i*) $\ker(\mathbf{L}_{1,\omega}^\lambda) = \{\mathbf{0}\}$.

(*ii*) $\ker(\mathbf{L}_{2,\omega}^\lambda) = \mathrm{span}\{\mathbf{\Phi}_0^\lambda\}$.

(*iii*) $n(\mathbf{L}_{1,\omega}^\lambda) < \infty$.

(*iv*) *The rest of the spectrum of the operators $\mathbf{L}_{1,\omega}^\lambda$ and $\mathbf{L}_{2,\omega}^\lambda$ is positive and bounded away from zero.*

Then the standing wave $e^{i\omega t}\mathbf{\Phi}_0^\lambda$ *is stable in* $H^1(\mathcal{G})$ *if* $n(\mathbf{L}_{1,\omega}^\lambda) - p(\omega_0) = 0$, *and* $e^{i\omega t}\mathbf{\Phi}_0^\lambda$ *is unstable if* $n(\mathbf{L}_{1,\omega}^\lambda) - p(\omega_0)$ *is odd.*

Below we give the description of the spectrum of the operators $\mathbf{L}_{1,\omega}^\lambda$ and $\mathbf{L}_{2,\omega}^\lambda$.

Proposition 4.4. *Let* $\lambda < 0$ *and* $\omega > \frac{N^2}{\lambda^2}$, *then the following assertions hold.*

(*i*) $\ker(\mathbf{L}_{2,\omega}^\lambda) = \operatorname{span}\{\mathbf{\Phi}_0^\lambda\}$ *and* $\mathbf{L}_{2,\omega}^\lambda \geq 0$.

(*ii*) *If* $\omega < \frac{N^2}{\lambda^2}\frac{p+1}{p-1}$, *then* $\ker(\mathbf{L}_{1,\omega}^\lambda) = \{\mathbf{0}\}$ *and* $n(\mathbf{L}_{1,\omega}^\lambda) = 1$.

(*iii*) *If* $\omega = \frac{N^2}{\lambda^2}\frac{p+1}{p-1}$, *then* $n(\mathbf{L}_{1,\omega}^\lambda) = 1$ *and* $\ker(\mathbf{L}_{1,\omega}^\lambda) = \operatorname{span}\{\mathbf{\Psi}_1^\lambda, .., \mathbf{\Psi}_{N-1}^\lambda\}$, *where*

$$\mathbf{\Psi}_j^\lambda = (0, .., 0, \underset{\mathbf{j}}{(\varphi_0^\lambda)'}, \underset{\mathbf{j+1}}{-(\varphi_0^\lambda)'}, 0, .., 0). \tag{4.5}$$

(*iv*) *If* $\omega > \frac{N^2}{\lambda^2}\frac{p+1}{p-1}$, *then* $\ker(\mathbf{L}_{1,\omega}^\lambda) = \{\mathbf{0}\}$ *and* $n(\mathbf{L}_{1,\omega}^\lambda) \leq N$. *Moreover, for* N *even in the space* $H^1_{\text{ev}}(\mathcal{G})$ *we have* $n(\mathbf{L}_{1,\omega}^\lambda) = 2$.

(*v*) *The rest of the spectrum of* $\mathbf{L}_{1,\omega}^\lambda$ *and* $\mathbf{L}_{2,\omega}^\lambda$ *is positive and bounded away from zero.*

Proof. (*i*) It is clear that $\mathbf{\Phi}_0^\lambda \in \ker(\mathbf{L}_{2,\omega}^\lambda)$. To show the equality $\ker(\mathbf{L}_{2,\omega}^\lambda) = \operatorname{span}\{\mathbf{\Phi}_0^\lambda\}$ let us note that any $\mathbf{V} = (v_j)_{j=1}^N \in H^2(\mathcal{G})$ satisfies the following identity

$$-v_j'' + \omega v_j - (\varphi_0^\lambda)^{p-1} v_j = \frac{-1}{\varphi_0^\lambda}\frac{d}{dx}\left[(\varphi_0^\lambda)^2 \frac{d}{dx}\left(\frac{v_j}{\varphi_0^\lambda}\right)\right], \quad x \neq 0.$$

Thus, for $\mathbf{V} \in D_\lambda$ we obtain

$$\begin{aligned}(\mathbf{L}_{2,\omega}^\lambda \mathbf{V}, \mathbf{V}) &= \sum_{j=1}^N \int_0^\infty (\varphi_0^\lambda)^2 \left(\frac{d}{dx}\left(\frac{v_j}{\varphi_0^\lambda}\right)\right)^2 dx + \sum_{j=1}^N \left[-v_j' v_j + v_j^2 \frac{(\varphi_0^\lambda)'}{\varphi_0^\lambda}\right]_0^\infty \\ &= \sum_{j=1}^N \int_0^\infty (\varphi_0^\lambda)^2 \left(\frac{d}{dx}\left(\frac{v_j}{\varphi_0^\lambda}\right)\right)^2 dx + \sum_{j=1}^N \left[v_j'(0) v_j(0) - v_j^2(0)\frac{(\varphi_0^\lambda)'(0)}{\varphi_0^\lambda(0)}\right].\end{aligned}$$

Due to the boundary conditions in (2.12) and formula (2.20) we obtain

$$\begin{aligned}\sum_{j=1}^N \left[v_j'(0) v_j(0) - v_j^2(0)\frac{(\varphi_0^\lambda)'(0)}{\varphi_0^\lambda(0)}\right] &= v_1'(0)\sum_{j=1}^N v_j(0) - \frac{N}{\lambda}\sum_{j=1}^N v_j^2(0) \\ &= \frac{1}{\lambda}\left(\left(\sum_{j=1}^N v_j(0)\right)^2 - N\sum_{j=1}^N v_j^2(0)\right),\end{aligned}$$

which is obviously positive for $\lambda < 0$ due to Jensen's inequality for the function $f(x) = x^2$.

Thus, $(\mathbf{L}_{2,\omega}^\lambda \mathbf{V}, \mathbf{V}) > 0$ for $\mathbf{V} \in D_\lambda \setminus \{\mathbf{\Phi}_0^\lambda, \mathbf{0}\}$ which proves (*i*).

(ii) Concerning the kernel of $\mathbf{L}^\lambda_{1,\omega}$, we recall that the only $L^2(\mathbb{R}_+)$-solution of the equation

$$-v_j'' + \omega v_j - p(\varphi_0^\lambda)^{p-1}v_j = 0$$

is $v_j = (\varphi_0^\lambda)'$ (up to a factor). Thus, any element of $\ker(\mathbf{L}^\lambda_{1,\omega})$ has the form $\mathbf{V} = (v_j)_{j=1}^N = (c_j(\varphi_0^\lambda)')_{j=1}^N$, $c_j \in \mathbb{C}$. If $v_1'(0) = ... = v_N'(0) \neq 0$, then by (2.12) we get $c_1 = ... = c_N \neq 0$ and consequently $N(\varphi_0^\lambda)'(0) = \lambda(\varphi_0^\lambda)''(0)$. Therefore, $\omega = \frac{N^2}{\lambda^2}$ which is impossible. Otherwise, the condition $v_j'(0) = 0$ implies that $(\varphi_0^\lambda)''(0) = 0$ which is equivalent to the identity $\omega = \frac{N^2}{\lambda^2}\frac{p+1}{p-1}$. Thus, we get that $c_1 = ... = c_j = 0$ and $\mathbf{V} \equiv \mathbf{0}$ for $\omega \neq \frac{N^2}{\lambda^2}\frac{p+1}{p-1}$.

The proof of the equality $n(\mathbf{L}^\lambda_{1,\omega}) = 1$ for $\omega < \frac{N^2}{\lambda^2}\frac{p+1}{p-1}$ is similar to the one in the previous case of NLS-δ. Namely, denoting

$$\boldsymbol{l}_\lambda = \left(\left(-\frac{d^2}{dx^2} + \omega - p(\varphi_0^\lambda)^{p-1}\right)\delta_{k,j}\right), \tag{4.6}$$

we define the following symmetric operator

$$\mathbf{L}_0' = \boldsymbol{l}_\lambda, \quad \mathrm{dom}(\mathbf{L}_0') = \left\{\mathbf{V} \in H^2(\mathcal{G}) : v_1'(0) = ... = v_N'(0) = 0,\ \sum_{j=1}^N v_j(0) = 0\right\}.$$

It is easily seen that $\mathbf{L}^\lambda_{1,\omega}$ is the self-adjoint extension of $\mathbf{L}_0'$. Let us show that the operator $\mathbf{L}_0'$ is non-negative. First, note that any $\mathbf{V} = (v_j)_{j=1}^N \in H^2(\mathcal{G})$ satisfies the following identity

$$-v_j'' + \omega v_j - p(\varphi_0^\lambda)^{p-1}v_j = \frac{-1}{(\varphi_0^\lambda)'}\frac{d}{dx}\left[((\varphi_0^\lambda)')^2\frac{d}{dx}\left(\frac{v_j}{(\varphi_0^\lambda)'}\right)\right], \quad x \neq 0.$$

Using the above equality and integrating by parts, we get for $\mathbf{V} \in \mathrm{dom}(\mathbf{L}_0')$

$$\begin{aligned}(\mathbf{L}_0'\mathbf{V}, \mathbf{V}) &= \sum_{j=1}^N \int_0^\infty ((\varphi_0^\lambda)')^2\left(\frac{d}{dx}\left(\frac{v_j}{(\varphi_0^\lambda)'}\right)\right)^2 dx + \sum_{j=1}^N \left[-v_j'v_j + v_j^2\frac{(\varphi_0^\lambda)''}{(\varphi_0^\lambda)'}\right]_0^\infty \\ &= \sum_{j=1}^N \int_0^\infty ((\varphi_0^\lambda)')^2\left(\frac{d}{dx}\left(\frac{v_j}{(\varphi_0^\lambda)'}\right)\right)^2 dx - \sum_{j=1}^N v_j^2(0)\frac{(\varphi_0^\lambda)''(0)}{(\varphi_0^\lambda)'(0)}.\end{aligned} \tag{4.7}$$

Taking into account that

$$-v_j^2(0)\frac{(\varphi_0^\lambda)''(0)}{(\varphi_0^\lambda)'(0)} = v_j^2(0)\frac{\lambda\omega}{2N}\left(p - 1 - (p+1)\frac{N^2}{\lambda^2\omega}\right),$$

we get nonnegativity of $\mathbf{L}_0'$ for $\omega \leq \frac{N^2}{\lambda^2}\frac{p+1}{p-1}$.

The adjoint operator is given by

$$(\mathbf{L}_0')^* = \boldsymbol{l}_\lambda, \quad \mathrm{dom}((\mathbf{L}_0')^*) = \left\{\mathbf{V} \in H^2(\mathcal{G}) : v_1'(0) = ... = v_N'(0)\right\}.$$

The last formula can be shown analogously to the previous case. Due to the Neumann decomposition, we get

$$\mathrm{dom}((\mathbf{L}_0')^*) = \big\{\mathbf{V} \in H^2(\mathcal{G}) : v_1'(0) = ... = v_N'(0)\big\} = \mathrm{dom}(\mathbf{L}_0') \dot{+} \mathrm{span}\{\mathbf{V}_i\} \dot{+} \mathrm{span}\{\mathbf{V}_{-i}\},$$

where $\mathbf{V}_{\pm i} = \Big(e^{i\sqrt{\pm i}x}\Big)_{j=1}^N$. Indeed, since $\varphi_0^\lambda \in L^\infty(\mathbb{R}_+)$, we get $\mathrm{dom}((\mathbf{L}_0')^*) = \mathrm{dom}(\widetilde{\mathbf{L}}^*)$, where

$$\widetilde{\mathbf{L}} = \Big(\Big(-\frac{d^2}{dx^2}\Big)\delta_{k,j}\Big), \quad \mathrm{dom}(\widetilde{\mathbf{L}}) = \mathrm{dom}(\mathbf{L}_0').$$

Since $\mathbf{L}_0' \geq 0$, deficiency indices are equal, and

$$n_\pm(\mathbf{L}_0') = \frac{\dim(\mathrm{dom}((\mathbf{L}_0')^*)/\,\mathrm{dom}(\mathbf{L}_0'))}{2} = 1.$$

By Proposition 3.4, $n(\mathbf{L}_{1,\omega}^\lambda) \leq 1$. Due to $(\mathbf{L}_{1,\omega}^\lambda \mathbf{\Phi}_0^\lambda, \mathbf{\Phi}_0^\lambda) = -(p-1)||\mathbf{\Phi}_0^\lambda||_{p+1}^{p+1} < 0$, we finally arrive at $n(\mathbf{L}_{1,\omega}^\lambda) = 1$, and *(ii)* is proved.

(iii) From the proof of item *(ii)* we induce that $n(\mathbf{L}_{1,\omega}^\lambda) = 1$ and the kernel of $\mathbf{L}_{1,\omega}^\lambda$ is nonempty as $\omega = \frac{N^2}{\lambda^2}\frac{p+1}{p-1}$. Moreover, we know that any element of the kernel has the form $\mathbf{V} = (v_j)_{j=1}^N = (c_j(\varphi_0^\lambda)')_{j=1}^N$, $c_j \in \mathbb{C}$, and it is necessary that $v_1'(0) = ... = v_N'(0) = 0$. Hence the condition

$$\lambda v_1'(0) = \sum_{j=1}^N v_j(0) = 0 \tag{4.8}$$

gives rise to $(N-1)$-dimensional kernel of $\mathbf{L}_{1,\omega}^\lambda$. Since the functions $\mathbf{\Psi}_j^\lambda$, $1 \leq j \leq N-1$, defined in (4.5) are linearly independent and satisfy the condition (4.8), they form the basis in $\ker(\mathbf{L}_{1,\omega}^\lambda)$, and *(iii)* is proven.

(iv) The identity $\ker(\mathbf{L}_{1,\omega}^\lambda) = \{\mathbf{0}\}$ was shown in *(ii)*. To show the inequality $n(\mathbf{L}_{1,\omega}^\lambda) \leq N$ we introduce the following minimal symmetric operator

$$\mathbf{L}_{\min} = \boldsymbol{l}_\lambda, \quad \mathrm{dom}(\mathbf{L}_{\min}) = \big\{\mathbf{V} \in H^2(\mathcal{G}) : v_1'(0) = ... = v_N'(0) = v_1(0) = ... = v_N(0) = 0\big\},$$

where $\boldsymbol{l}_\lambda$ is defined in (4.6). The operator $\mathbf{L}_{1,\omega}^\lambda$ is the self-adjoint extension of $\mathbf{L}_{\min}$. From the formula (4.7) it follows that $\mathbf{L}_{\min}$ is a non-negative operator. It is obvious that

$$\mathbf{L}_{\min}^* = \boldsymbol{l}_\lambda, \quad \mathrm{dom}(\mathbf{L}_{\min}^*) = H^2(\mathcal{G}).$$

Then, due to the Neumann formula

$$\mathrm{dom}(\mathbf{L}_{\min}^*) = \mathrm{dom}(\mathbf{L}_{\min}) \dot{+} \mathrm{span}\{\mathbf{V}_i^1, .., \mathbf{V}_i^N\} \dot{+} \mathrm{span}\{\mathbf{V}_{-i}^1, .., \mathbf{V}_{-i}^N\},$$

where $\mathbf{V}_{\pm i}^j = \Big(0, .., \underset{\mathbf{j}}{e^{i\sqrt{\pm i}x}}, 0, .., 0\Big)$, and consequently $n_\pm(\mathbf{L}_{\min}) = N$. By Proposition 3.4, $n(\mathbf{L}_{1,\omega}^\lambda) \leq N$.

Let N be even. It is easily seen that $n_\pm(\mathbf{L}_{\min}) = 2$ in $H_{\mathrm{ev}}^1(\mathcal{G})$. Indeed,

$$\mathrm{dom}(\mathbf{L}_{\min}^*) = \mathrm{dom}(\mathbf{L}_{\min}) \dot{+} \mathrm{span}\{\widetilde{\mathbf{V}}_i^1, \widetilde{\mathbf{V}}_i^2\} \dot{+} \mathrm{span}\{\widetilde{\mathbf{V}}_{-i}^1, \widetilde{\mathbf{V}}_{-i}^2\},$$

where

$$\widetilde{\mathbf{V}}^1_{\pm i} = (\underset{\mathbf{1}}{e^{i\sqrt{\pm i}x}}, ..., \underset{\mathbf{N/2}}{e^{i\sqrt{\pm i}x}}, \underset{\mathbf{N/2+1}}{0}, ..., \underset{\mathbf{N}}{0}), \quad \widetilde{\mathbf{V}}^2_{\pm i} = (\underset{\mathbf{1}}{0}, ..., \underset{\mathbf{N/2}}{0}, \underset{\mathbf{N/2+1}}{e^{i\sqrt{\pm i}x}}, ..., \underset{\mathbf{N}}{e^{i\sqrt{\pm i}x}}).$$

Thus, by Proposition 3.4, we get $n(\mathbf{L}^\lambda_{1,\omega}) \leq 2$ in $H^1_{\mathrm{ev}}(\mathcal{G})$.

Let us introduce the following quadratic form $F^\lambda_{1,\omega}$ associated with the operator $\mathbf{L}^\lambda_{1,\omega}$

$$\begin{aligned}&F^\lambda_{1,\omega}(\mathbf{V}) = ||\mathbf{V}'||^2 + \omega||\mathbf{V}||^2 - p\sum_{j=1}^{N}\int\limits_0^\infty (\varphi_0^\lambda)^{p-1} v_j^2 dx + \tfrac{1}{\lambda}\left(\sum_{j=1}^{N} v_j(0)\right)^2,\\ &\mathrm{dom}(F^\lambda_{1,\omega}) = H^1(\mathcal{G}).\end{aligned}$$

Consider

$$\widetilde{\mathbf{\Psi}} = (\underset{\mathbf{1}}{(\varphi_0^\lambda)'}, ..., \underset{\mathbf{N/2}}{(\varphi_0^\lambda)'}, \underset{\mathbf{N/2+1}}{-(\varphi_0^\lambda)'}, ..., \underset{\mathbf{N}}{-(\varphi_0^\lambda)'}).$$

Integrating by parts and using (4.5), we get

$$\begin{aligned}F^\lambda_{1,\omega}(\widetilde{\mathbf{\Psi}}) &= N\int\limits_0^\infty (\varphi_0^\lambda)'\left(-(\varphi_0^\lambda)''' + \omega(\varphi_0^\lambda)' - p(\varphi_0^\lambda)^{p-1}(\varphi_0^\lambda)'\right)dx - N(\varphi_0^\lambda)'(0)(\varphi_0^\lambda)''(0)\\ &= \tfrac{N^2}{2\lambda}\omega\left(\tfrac{(p+1)\omega}{2}\right)^{\frac{2}{p-1}}\left(1 - \tfrac{N^2}{\lambda^2\omega}\right)^{\frac{2}{p-1}}\left(p - 1 - (p+1)\tfrac{N^2}{\lambda^2\omega}\right),\end{aligned}$$

which is negative for $\omega > \frac{N^2}{\lambda^2}\frac{p+1}{p-1}$. Recalling that $(\mathbf{L}^\lambda_{1,\omega}\mathbf{\Phi}^\lambda_0, \mathbf{\Phi}^\lambda_0) < 0$, we get by orthogonality of $\widetilde{\mathbf{\Psi}}$ and $\mathbf{\Phi}^\lambda_0$

$$F^\lambda_{1,\omega}(s\mathbf{\Phi}^\lambda_0 + r\widetilde{\mathbf{\Psi}}) = |s|^2 F^\lambda_{1,\omega}(\mathbf{\Phi}^\lambda_0) + |r|^2 F^\lambda_{1,\omega}(\widetilde{\mathbf{\Psi}}) < 0, \quad r, s \in \mathbb{C}.$$

Thus, $F^\lambda_{1,\omega}$ is negative on two-dimensional subspace $\mathcal{M} = \mathrm{span}\{\mathbf{\Phi}^\lambda_0, \widetilde{\mathbf{\Psi}}\}$. Therefore, by Proposition 3.6, we get $n(\mathbf{L}^\lambda_{1,\omega}) \geq 2$. The assertion (iv) is proven. The proof of item (v) is standard and relies on the Weyl theorem. This finishes the proof of the Proposition.

□

Finally, we have to evaluate $p(\omega)$ defined in (4.4) estimating $\partial_\omega||\mathbf{\Phi}^\lambda_0||^2$.

Proposition 4.5. *Let* $\omega > \frac{N^2}{\lambda^2}$ *and* $J(\omega) = \partial_\omega||\mathbf{\Phi}^\lambda_0||^2$.

(i) *If* $1 < p \leq 5$, *then* $J(\omega) > 0$.

(ii) *If* $p > 5$, *then there exists* ω^* *such that* $J(\omega^*) = 0$, *and* $J(\omega) > 0$ *for* $\omega \in \left(\frac{N^2}{\lambda^2}, \omega^*\right)$, *while* $J(\omega) < 0$ *for* $\omega \in (\omega^*, \infty)$.

Proof. Recall that $\mathbf{\Phi}_0^\lambda=(\varphi_0^\lambda)_{j=1}^N$, where φ_0^λ is defined by (2.20). Changing variables we get

$$\int\limits_0^\infty(\varphi_0^\lambda(x))^2dx=\left(\frac{p+1}{2}\right)^{\frac{2}{p-1}}\frac{2\omega^{\frac{2}{p-1}-\frac12}}{p-1}\int\limits_{\tanh^{-1}\left(\frac{N}{|\lambda|\sqrt{\omega}}\right)}^\infty\operatorname{sech}^{\frac{4}{p-1}}y\,dy$$

$$=\left(\frac{p+1}{2}\right)^{\frac{2}{p-1}}\frac{2\omega^{\frac{2}{p-1}-\frac12}}{p-1}\int\limits_{\frac{N}{|\lambda|\sqrt{\omega}}}^1(1-t^2)^{\frac{2}{p-1}-1}dt.$$

From the last equality we get

$$J(\omega)=C\omega^{\frac{7-3p}{2(p-1)}}\left[\tfrac{5-p}{p-1}\int\limits_{\frac{N}{|\lambda|\sqrt{\omega}}}^1(1-t^2)^{\frac{3-p}{p-1}}dt+\tfrac{N}{|\lambda|\sqrt{\omega}}(1-\tfrac{N^2}{\lambda^2\omega})^{\frac{3-p}{p-1}}\right]=C\omega^{\frac{7-3p}{2(p-1)}}J_1(\omega),\qquad(4.9)$$

where $C=\frac{N}{p-1}\left(\frac{p+1}{2}\right)^{\frac{2}{p-1}}>0$ and

$$J_1(\omega)=\left[\tfrac{5-p}{p-1}\int\limits_{\frac{N}{|\lambda|\sqrt{\omega}}}^1(1-t^2)^{\frac{3-p}{p-1}}dt+\tfrac{N}{|\lambda|\sqrt{\omega}}(1-\tfrac{N^2}{\lambda^2\omega})^{\frac{3-p}{p-1}}\right].$$

Thus,

$$J_1'(\omega)=\tfrac{N}{|\lambda|\omega^{3/2}}\tfrac{3-p}{p-1}\left[\left(1-\tfrac{N^2}{\lambda^2\omega}\right)^{\frac{3-p}{p-1}}+\tfrac{N^2}{\lambda^2\omega}\left(1-\tfrac{N^2}{\lambda^2\omega}\right)^{-\frac{2(p-2)}{p-1}}\right].\qquad(4.10)$$

It is immediate that $J(\omega)>0$ for $1<p\leq5$. Consider the case $p>5$. It is easily seen that $J_1(\omega)>0$ for ω in a neighborhood of $\frac{N^2}{\lambda^2}$ and $J_1(\omega)<0$ for ω big enough. Moreover, from (4.10) it follows that $J_1'(\omega)<0$ for $\omega>\frac{N^2}{\lambda^2}$ and consequently $J_1(\omega)$ is strictly decreasing. Therefore, there exists a unique $\omega^*>\frac{N^2}{\lambda^2}$ such that $J_1(\omega^*)=J(\omega^*)=0$, consequently $J(\omega)>0$ for $\omega\in\left(\frac{N^2}{\lambda^2},\omega^*\right)$ and $J(\omega)<0$ for $\omega\in(\omega^*,\infty)$. □

Proof of Theorem 1.1

- (*i*) Combining Theorem 4.3, Proposition 4.4 (items (*i*), (*ii*) and (*v*)), and Proposition 4.5 (*i*) we get assertion (*i*).
- (*ii*) Combining Theorem 4.3, Proposition 4.4 (items (*i*), (*iv*) and (*v*)), and Proposition 4.5 (*i*) we get instability of $e^{i\omega t}\mathbf{\Phi}_0^\lambda$ (*i*) in $H^1_{\rm ev}(\mathcal{G})$. We note that well-posedness of equation (2.11) in $H^1_{\rm ev}(\mathcal{G})$ follows from the uniqueness of the solution to the Cauchy problem. Finally, instability in the smaller space $H^1_{\rm ev}(\mathcal{G})$ induces instability in all $H^1(\mathcal{G})$.

□

Remark 4.6. (*i*) Relative position of ω^* and $\frac{N^2}{\lambda^2}\frac{p+1}{p-1}$ is not clear. Computer analysis gives us an opportunity to claim the hypothesis that $\omega^* > \frac{N^2}{\lambda^2}\frac{p+1}{p-1}$. The last one by Theorem 4.3 implies the following hypothesis on the stability of $\mathbf{U}(t,x) = e^{i\omega t}\mathbf{\Phi}_0^\lambda$ defined in (2.20) in the case $p > 5$:

(*a*) If $\omega \in \left(\frac{N^2}{\lambda^2}, \frac{N^2}{\lambda^2}\frac{p+1}{p-1}\right)$, then $\mathbf{U}(t,x)$ is stable in $H^1(\mathcal{G})$.

(*b*) If $\omega \in \left(\frac{N^2}{\lambda^2}\frac{p+1}{p-1}, \omega^*\right)$ and N is even, then $\mathbf{U}(t,x)$ is unstable in $H^1(\mathcal{G})$.

(*c*) If $\omega > \omega^*$ and N is odd, then $\mathbf{U}(t,x)$ is unstable in $H^1(\mathcal{G})$.

(*ii*) In the case $\omega = \omega^*$ and $p > 5$ we conjecture due to [35, Corollary 2] that the standing wave $\mathbf{U}(t,x)$ is unstable. Indeed, it is easy to show that the number of negative eigenvalues of the operator

$$\widetilde{\mathbf{L}}_{1,\omega}^\lambda = \left(\left(-\frac{d^2}{dx^2} + \omega - p(\varphi_0^\lambda)^{p-1}\right)\delta_{k,j}\right) \qquad \mathrm{dom}(\widetilde{\mathbf{L}}_{1,\omega}^\lambda) = D_\lambda \cap \mathcal{E}(\mathcal{G}),$$

equals one. Let $c(\omega) = C\omega^{\frac{7-3p}{2(p-1)}}$, then from (4.9) we get

$$J(\omega) = c(\omega)J_1(\omega).$$

Therefore using the properties of $J_1(\omega)$ established at the proof of Proposition 4.5, we get

$$J'(\omega^*) = c'(\omega^*)J_1(\omega^*) + c(\omega^*)J_1'(\omega^*) = c(\omega^*)J_1'(\omega^*) \neq 0,$$

or $\partial_\omega^2||\mathbf{\Phi}_0^\lambda||^2(\omega^*) \neq 0$. Thus, by [35, Corollary 2] we can conjecture that for $\omega = \omega^*$ the standing wave $\mathbf{U}(t,x)$ is unstable in $\mathcal{E}(\mathcal{G})$ and consequently in $H^1(\mathcal{G})$.

Acknowledgements.

J. Angulo was supported partially by CNPq/Brazil, N. Goloshchapova was supported by FAPESP under the projects 2012/50503-6 and 2016/02060-9.

References

[1] R. Adami, C. Cacciapuoti, D. Finco, D. Noja, Stable standing waves for a NLS on star graphs as local minimizers of the constrained energy, arXiv:1509.01810v1.

[2] R. Adami, C. Cacciapuoti, D. Finco, D. Noja, Variational properties and orbital stability of standing waves for NLS equation on a star graph, J. Differential Equations **257** (2014), no. 10, 3738–3777.

[3] R. Adami and D. Noja, Stability and symmetry-breaking bifurcation for the ground states of a NLS with a δ' interaction, Comm. Math. Phys. **318** (2013), no. 1, 247–289.

[4] R. Adami, D. Noja and N. Visciglia, Constrained energy minimization and ground states for NLS with point defects, Discrete Contin. Dyn. Syst. Ser. B **18** (2013), no. 5, 1155–1188.

[5] R. Adami and D. Noja, Existence of dynamics for a 1D NLS equation perturbed with a generalized point defect, J. Phys. A **42** (2009), no. 49, 495302, 19 pp.

[6] G. Agrawal, *Nonlinear Fiber Optic*, Academic Press, 2001.

[7] S. Albeverio, F. Gesztesy, R. Hoegh-Krohn, H. Holden, *Solvable models in quantum mechanics*, second edition, AMS Chelsea Publishing, Providence, RI, 2005.

[8] S. Albeverio and P. Kurasov, *Singular perturbations of differential operators*, London Mathematical Society Lecture Note Series, 271, Cambridge Univ. Press, Cambridge, 2000.

[9] W.O. Amrein, A.M. Hinz, D.P. Pearson, *Sturm-Liouville theory*, Birkhäuser Verlag, Basel, 2005.

[10] J. Angulo Pava, Instability of cnoidal-peak for the NLS-δ-equation, Math. Nachr. **285** (2012), no. 13, 1572–1602.

[11] J. Angulo Pava, N. Goloshchapova, Stability of standing waves for NLS-log equation with δ-interaction, arXiv:1506.08455v2.

[12] J. Angulo Pava and G. Ponce, The non-linear Schrödinger equation with a periodic δ-interaction, Bull. Braz. Math. Soc. (N.S.) **44** (2013), no. 3, 497–551.

[13] G. Berkolaiko and P. Kuchment, *Introduction to quantum graphs*, Mathematical Surveys and Monographs, 186, Amer. Math. Soc., Providence, RI, 2013.

[14] H. Berestycki and T. Cazenave, Instabilité des états stationnaires dans les équations de Schrödinger et de Klein-Gordon non linéaires, C. R. Acad. Sci. Paris Sér. I Math. **293** (1981), no. 9, 489–492.

[15] F. A. Berezin and M. A. Shubin, *The Schrödinger equation*, translated from the 1983 Russian edition by Yu. Rajabov, D. A. Leĭtes and N. A. Sakharova and revised by Shubin, Mathematics and its Applications (Soviet Series), 66, Kluwer Acad. Publ., Dordrecht, 1991.

[16] J. Blank, P. Exner, M. Havlicek, *Hilbert space operators in quantum physics*, second edition, Theoretical and Mathematical Physics, Springer, New York, 2008.

[17] V.A. Brazhnyi, V.V. Konotop, Theory of nonlinear matter waves in optical lattices, Mod. Phys. Lett. B **18** (2004), 627–551.

[18] X. D. Cao and B. A. Malomed, Soliton-defect collisions in the nonlinear Schrödinger equation, Phys. Lett. A **206** (1995), no. 3-4, 177–182.

[19] V. Caudrelier, M. Mintchev and E. Ragoucy, Solving the quantum nonlinear Schrödinger equation with δ-type impurity, J. Math. Phys. **46** (2005), no. 4, 042703, 24 pp.

[20] T. Cazenave and P.-L. Lions, Orbital stability of standing waves for some nonlinear Schrödinger equations, Comm. Math. Phys. **85** (1982), no. 4, 549–561.

[21] T. Cazenave, *An Introduction to Nonlinear Schrödinger Equations*, Textos de Métodos Mathematicos, **26**, IM-UFRJ, Rio de Janeiro, 1989.

[22] S. Le Coz, R. Fukuizumi, G. Fibich, B. Ksherim, Y. Sivan, Instability of bound states of a nonlinear Schrödinger equation with a Dirac potential, Phys. D **237** (2008), no. 8, 1103–1128.

[23] R. Fukuizumi and L. Jeanjean, Stability of standing waves for a nonlinear Schrödinger equation with a repulsive Dirac delta potential, Discrete Contin. Dyn. Syst. **21** (2008), no. 1, 121–136.

[24] R. Fukuizumi, M. Ohta and T. Ozawa, Nonlinear Schrödinger equation with a point defect, Ann. Inst. H. Poincaré Anal. Non Linéaire **25** (2008), no. 5, 837–845.

[25] R. H. Goodman, P. J. Holmes and M. I. Weinstein, Strong NLS soliton-defect interactions, Phys. D **192** (2004), no. 3-4, 215–248.

[26] M. Grillakis, J. Shatah and W. Strauss, Stability theory of solitary waves in the presence of symmetry. I, J. Funct. Anal. **74** (1987), no. 1, 160–197.

[27] M. Grillakis, J. Shatah and W. Strauss, Stability theory of solitary waves in the presence of symmetry. II, J. Funct. Anal. **94** (1990), no. 2, 308–348.

[28] J. Holmer, J. Marzuola and M. Zworski, Fast soliton scattering by delta impurities, Comm. Math. Phys. **274** (2007), no. 1, 187–216.

[29] J. Holmer, J. Marzuola and M. Zworski, Soliton splitting by external delta potentials, J. Nonlinear Sci. **17** (2007), no. 4, 349–367.

[30] M. Kaminaga and M. Ohta, Stability of standing waves for nonlinear Schrödinger equation with attractive delta potential and repulsive nonlinearity, Saitama Math. J. **26** (2009), 39–48.

[31] P. Kuchment, Quantum graphs. I. Some basic structures, Waves Random Media **14** (2004), no. 1, S107–S128.

[32] C.R. Menyuk, Soliton robustness in optical fibers, J. Opt. Soc. Am. B **10** (1993), no. 9, 1585–1591.

[33] J. Moloney, A. Newell, *Nonlinear optics*, Advanced Topics in the Interdisciplinary Mathematical Sciences, Addison-Wesley, Redwood City, CA, 1992.

[34] M.A. Naimark, *Linear differential operators* (Russian), second edition, revised and augmented., Izdat. "Nauka", Moscow, 1969.

[35] M. Ohta, Instability of bound states for abstract nonlinear Schrödinger equations, J. Funct. Anal. **261** (2011), no. 1, 90–110.

[36] M. Reed and B. Simon, *Methods of modern mathematical physics. IV. Analysis of operators*, Academic Press, New York, 1978.

[37] H. Sakaguchi and M. Tamura, Scattering and trapping of nonlinear Schrödinger solitons in external potentials, J. Phys. Soc. Japan **73** (2004), no. 3, 503–506.

[38] B.T. Seaman, L.D. Carr, M.J. Holland, Effect of a potential step or impurity on the Bose-Einstein condensate mean field, Phys. Rev. A **71** (2005), 033602.

[39] M. I. Weinstein, Nonlinear Schrödinger equations and sharp interpolation estimates, Comm. Math. Phys. **87** (1982/83), no. 4, 567–576.

Jaime Angulo Pava
Department of Mathematics, IME-USP,
Rua do Matão 1010, Cidade Universitária,
CEP 05508-090, São Paulo, SP, Brazil
e-mail: angulo@ime.usp.br

Nataly Goloshchapova,
Department of Mathematics, IME-USP,
Rua do Matão 1010, Cidade Universitária,
CEP 05508-090, São Paulo, SP, Brazil
e-mail: nataliia@ime.usp.br